

The Higgs sector of gravitational gauge theories

M. Leclerc

*Section of Astrophysics and Astronomy, Department of Physics,
University of Athens, Greece*

Abstract

Gravitational gauge theories with de Sitter, Poincaré and affine symmetry group are investigated under the aspect of the breakdown of the initial symmetry group down to the Lorentz subgroup. As opposed to the nonlinear realization approach, in the dynamical symmetry breaking procedure, the structure subgroup is not chosen arbitrarily, but is dictated by the symmetry of the groundstate of a Higgs field. We review the theory of spontaneously broken de Sitter gravity by Stelle and West and apply a similar approach to the case of the Poincaré and affine groups. We will find that the Poincaré case is almost trivial. The translational Higgs field reveals itself as pure gauge, i.e., it is expressed entirely in terms of the Nambu-Goldstone bosons and does not appear in the Lagrangian after the symmetry breaking. The same holds for the translational part of the affine group. The Higgs field provoking the breakdown of the general linear group leads to the determination of the Lorentzian signature of the metric in the groundstate. We show that the Higgs field remains in its groundstate, i.e., that the metric will have Lorentzian signature, unless we introduce matter fields that explicitly couple to the symmetric part of the connection. Furthermore, we present arguments that the Lorentzian signature is actually the only possible choice for physical spacetime, since the symmetry breaking mechanism works only if the stability subgroup is taken to be the Lorentz group. The other four-dimensional rotation groups are therefore ruled out not only on physical, but also on theoretical grounds. Finally, we show that some features, like the necessity of the introduction of a dilaton field, that seem artificial in the context of the affine theory, appear most natural if the gauge group is taken to be the special linear group in five dimensions. We also present an alternative model which is based on the spinor representation of the Lorentz group and is especially adopted to the description of spinor fields in a general linear covariant way, without the use of the infinite dimensional representations which are usually considered to be unavoidable.

PACS: 04.50.+h; 11.15.Ex

1 Introduction

That classical gravity represents the subsidiary of a theory with larger symmetry has been suggested in the past, presenting the theory in terms of nonlinear realizations of those groups. This procedure, however, requires that the Lorentz subgroup is chosen arbitrarily as structure group and therefore, does not explain the Minkowskian structure of the physical spacetime. On the other hand, introducing explicitly a Higgs sector into the theory, results in a dynamical symmetry breaking, with a groundstate invariant under the Lorentz subgroup only. Such a model has been presented by Stelle and West in the case of the de Sitter group. We will review their results and apply a similar approach to the Poincaré and affine groups.

The concept of nonlinear realizations has been introduced shortly after the advent of spontaneously broken theories [1, 2, 3] and its most successful candidate, the electroweak theory of Weinberg and Salam [4], by Coleman, Wess and Zumino in the context of gauge theories of internal symmetry groups [5] (see also [6]). Applying these concepts to the general linear group $GL(R^4)$, Isham, Salam and Strathdee [7] formulated the first version of a *broken* gauge theory of gravity. Since then, many attempts have been made to use nonlinear realizations in the context of gravity, see for instance [8, 9, 10] for the (super)Poincaré group, [11] for the de Sitter group, [12] for the affine group and [13, 14] for attempts to include diffeomorphism groups. The conformal group has also received considerable attention [7, 13, 15, 16]. Other references can be found in the cited articles.

In the nonlinear realization approach, the structure (or stability) subgroup H of the symmetry group G can be chosen arbitrarily. This is a result of the fact that the choice of a certain subgroup essentially consists in parameterizing some of the gauge degrees of freedom. Simply stated, some parts of the connection are transformed into tensors with the help of Nambu-Goldstone fields and the corresponding gauge freedom is transferred to those fields. This is not a gauge choice or gauge fixing, but rather a gauge parameterization. As such, it does not effect the G symmetry of the theory in any way, and therefore the choice of one or another subgroup is necessarily equivalent.

Clearly, this is somehow unsatisfactory, since we know from everyday physics that the residual symmetry group should be the Lorentz group $SO(3,1)$ (or eventually the Poincaré group) and not, say, the de Sitter group or the group $SO(4)$. After all, there is little doubt that we live in a 4-dimensional Minkowskian world and not in a 5-dimensional or Euclidian one. It is therefore natural to look for a mechanism

that fixes the choice of the structure group as a result of the dynamics of the theory. The introduction of a Higgs field, carrying a representation of the symmetry group G , with a suitable Higgs potential leading to a groundstate that is not symmetric under G anymore, but only under the Lorentz subgroup $H = SO(3, 1)$ provides such a mechanism.

That gravity is a dynamically broken gauge theory has been suggested in the past, but with the exception of the work of Stelle and West [11], we have never really seen neither a kinetic term for the Higgs field, nor a potential provoking the symmetry breakdown. We refer here only to theories that are constructed, as far as possible, without the help of additional ingredients, merely in terms of the connection of the symmetry group and those fields that are unavoidable to write down a meaningful Lagrangian density, as are the Poincaré coordinates or the metric in the case of the general linear group.

It is most astonishing, especially in the case of Poincaré gauge theory, which has raised the interest of so many people, that although the introduction of the Nambu-Goldstone field (also called Poincaré coordinates, or Cartan's radius vector) in the manner of [10, 17] and [18] (whose ultimate interpretation is provided by the nonlinear realization approach, see [10, 19]) is widely accepted, nobody seems to have presented the Higgs sector that is supposed to break down the symmetry from the Poincaré to the Lorentz group. Let us cite a characteristic statement from the standard reference on metric affine gauge theory by Hehl et al. [20]: *We believe that the story of the ξ (the Poincaré coordinates) has not yet come to an end and that future developments on this point are possible. Probably, one has to come up with an idea of how to construct an explicit symmetry breaking mechanism.* We will show that, depending on the degree of optimism in the physicist's viewpoint, the Higgs sector for this breaking mechanism is actually already implicitly incorporated in the standard Poincaré gauge theory framework, or, taking a less optimistic viewpoint, that no Higgs sector (other than that) can be constructed.

On the other hand, symmetry breaking mechanisms have been presented in the case of the general linear group. However, the Higgs sector in those cases is constructed with the help of an additional field in a general linear, infinite dimensional representation (a so called manifold) [12, 21]. This is unsatisfying from several points of view. First, it does not explain the role of the general linear metric tensor, which is a necessary ingredient of the theory since no Lagrangian invariant under the general linear group can be written down without the help of a metric. The nonlinear realization approach

clearly suggests that it is this tensor that should play the role of the Higgs field and trigger the symmetry breaking down to the Lorentz group. This avoids not only the use of infinite dimensional representations, but also the introduction of yet another field to the already complicated structure of the theory. Secondly, it is not really clear to us how the manifold approach favors the Lorentz group as stability subgroup as opposed to the other possible rotation groups, like $O(4)$ etc. We will present a much simpler approach, with the metric playing the role of the Higgs field, and the symmetry breaking fixing the signature of the groundstate metric.

However, it turns out, that this method actually works only in the case where the symmetry group is taken to be the special linear group. In order to reduce the general linear group to the special linear group, we will have to introduce an additional Higgs dilaton field, following a similar approach as in [20]. We present arguments that explain why the introduction of this field (which is not really apparent in the nonlinear realization scheme) is unavoidable. In brief, the reason can be traced back to the fact that, in the metric affine theory, one of the field equations is redundant. Moreover, we will show that the dilaton approach contains a small loophole, and in order to construct a consistent Higgs sector, an additional scalar field has to be included.

It turns out that the introduction of both the scalar and the dilaton field can be naturally explained in the context of the gauge theory of the special linear group $SL(R^5)$. This theory combines the features of the affine and de Sitter theories and provides us with a mechanism to break down the symmetry to the Lorentz group without any additional fields.

In order to avoid misunderstandings, it is important to point out that throughout the article, the gauge group is treated exactly in the same way as in conventional gauge theories of internal symmetries. Thus, the gauge potential is a connection one-form transforming under the group G in the usual way (see equ. (4) below), this gauge group being in no way related to spacetime coordinate transformations. The difference between our theories and conventional Yang-Mills theories will arise through additional structure, like tetrad fields or metrics, that will be defined with the help of the connection and of the Higgs fields, a posteriori. This conception of gauge gravity coincides essentially with the one described in [19] and [20]. It seems to us the most promising way in view of a possible unification of gravity with the other gauge interactions. It is, however, not the only way. Alternative theories, based on a different gauge concept, have been presented and analyzed in detail in the past. In those theories, the general linear gauge group is directly related to the coordinate

transformations and the tetrad fields arise as the Goldstone fields of the symmetry breakdown. In other words, the quotient space G/H , where H is the Lorentz group, is identified with physical spacetime, and not, as in our approach, with a tangent space to the spacetime manifold. We will not deal with such theories in this article and refer the reader to the original article of Ivanenko and Sardanashvily [22] as well as to more recent work, e.g., [23, 24]. The later, dealing with the coupling of gravity to spinor matter, is essentially interesting in connection with section 7 of this article. Note also that the approach of [12, 13, 14], and also of one of the earliest Higgs approaches to gravity by Borisov and Ogievetsky [25], are based on a similar gauge concept concerning the general linear group. Without going into details, it is clear what are the weak and strong points of both, alternative, viewpoints. As outlined above, the approach of Hehl et al., which we adopt here, is more suitable in view of a unification of the fundamental forces, since, in a sense, it is based on the idea to describe gravity in a way as close as possible to the description of the other forces of nature. On the other hand, the diffeomorphism approach of [22] essentially starts by underlining not the similarities to other interactions, but rather the features that are unique to gravity, namely the equivalence principle and the universality, which are then married with a suitable gauge concept. This is certainly a promising way too, since from those same principles, Einstein was led successfully to general relativity in the first place. In our approach, the validity of those principles has to be established a posteriori, while the fundamental concept is the gauge principle. This is not necessarily a drawback, since it is not evident whether the equivalence principle is valid in generalized gravity theories, or whether it represents just a sort of a classical, macroscopic limit. For instance, in Poincaré gauge theory, as is well known, the equivalence principle is certainly not generally valid, since particles with different intrinsic spin follow different trajectories [26]. Interesting remarks concerning the different gauge concepts are also found in [27].

We will begin this article by reviewing the gauge theory of the de Sitter group as it was presented by Stelle and West twenty five years ago. This gives us the possibility to follow step by step the same method in the other cases, which present some particularities, and which else would hardly be recognizable as dynamically broken theories.

In detail, the article is organized as follows. After a short excursion to the nonlinear realization approach (section 2), we will take the work of Stelle and West as a starting point for the construction of the Higgs sector of gauge theories of gravity for different symmetry groups. We start by reviewing the de Sitter case (section 3) and go on to the

Poincaré and affine groups in sections 4 and 5. Finally, a short analysis of the $SL(R^5)$ theory will be given in section 6, focusing on the similarities to the affine case. In a last section, we briefly describe an alternative Higgs mechanism where the general linear symmetry is broken down to the group $SL(2, C)$, i.e., to the spinor representation of the Lorentz group.

2 Nonlinear realizations

We very briefly review the basic concepts of this approach, in order to compare later on with the conventional Higgs mechanism. A modern and compact description of the method can be found in [19], in the context of the affine group.

Let G be the symmetry group of the theory and H the structure subgroup (which is chosen arbitrarily in this approach). Consider a section on the principal fiber bundle $G(G/H, H)$,

$$\sigma : G/H \rightarrow G. \quad (1)$$

Then, any element g of G can be written uniquely as

$$g = \sigma(\xi)h, \quad (2)$$

with $\xi \in G/H$ and $h \in H$. Especially, the element $g\sigma(\xi)$ can be written as

$$g\sigma(\xi) = \sigma(\xi')h, \quad (3)$$

for some $h \in H$. Obviously, the coset parameters ξ' as well as h will depend on ξ and g , i.e., $\xi' = \xi'(\xi, g)$ and $h = h(\xi, g)$. A short form of (3) is $\sigma' = g\sigma h^{-1}$.

If we have a linear connection transforming under G as

$$\Gamma \rightarrow g\Gamma g^{-1} + g dg^{-1}, \quad (4)$$

then the following quantity

$$\tilde{\Gamma} = \sigma^{-1}\Gamma\sigma + \sigma^{-1}d\sigma, \quad (5)$$

clearly transforms as

$$\tilde{\Gamma} \rightarrow h\tilde{\Gamma}h^{-1} + h dh^{-1}, \quad (6)$$

i.e., basically as a (nonlinear) connection under the group H (apart from those components that are outside of the Lie algebra of H , which transform as tensors under H , see [19]).

More generally, to any field φ transforming under some representation of G as $\varphi \rightarrow g\varphi$, we can associate a field $\psi = \sigma^{-1}\varphi$ that will transform as $\psi \rightarrow h\psi$. This presupposes of course that the transformation law $\sigma \rightarrow g\sigma h^{-1}$ is interpreted as transformation under the corresponding representations. We will come back to this important point in section 7.

Let us write the abstract formulas (3) to (6) in component form. Quantities transforming under the group G will be labeled by greek indices $\alpha, \beta \dots$ and those transforming under the subgroup H by latin letters from the beginning of the alphabet $a, b, c, \dots$ (We reserve latin letters from the middle of the alphabet $i, j, k \dots$ for spacetime indices.) Equation (3) in the form $\sigma' = g\sigma h^{-1}$ shows that σ transforms as a mixed tensor. We will denote it by r^α_a and its inverse by r^α_α . Let Γ^α_β be the G -connection 1-form, i.e., we have the transformation law

$$\Gamma^\alpha_\beta \rightarrow G^\alpha_\gamma \Gamma^\gamma_\delta (G^{-1})^\delta_\beta + G^\alpha_\gamma d(G^{-1})^\gamma_\beta, \quad (7)$$

where G^α_β is an element of G . Equation (5) defines the H -connection as

$$\Gamma^a_b = r^\alpha_a \Gamma^\alpha_\beta r^\beta_b + r^\alpha_a dr^\alpha_b. \quad (8)$$

Using the transformation law for r^α_a

$$r^\alpha_a \rightarrow G^\alpha_\beta r^\beta_b (H^{-1})^b_a, \quad (9)$$

with $H^a_b \in H$, we find

$$\Gamma^a_b \rightarrow \Gamma^a_b = H^a_c \Gamma^c_d (H^{-1})^d_b + H^a_c d(H^{-1})^c_b, \quad (10)$$

which is equation (6) again. Thus, we have reduced the G -connection down to a H -connection (plus tensor parts). The matrix r^α_b is also called *reducing matrix* [6]. Note that you may see (8) as a gauge transformation (it is of the same form as (7), with r^α_b as a special element of G). Thus, in a sense, the reduction (8) is a gauge choice, but the gauge freedom is not lost, the degrees of freedom have just been transferred to the reducing matrix. That is what we meant in the introduction by *gauge parameterization*.

Let us close this section by demonstrating this formalism with an explicit example. For simplicity, we take the Poincaré group $ISO(3, 1)$ as symmetry group G which reduces to the Lorentz subgroup $H = SO(3, 1)$. How can we find a suitable reducing matrix? Well, as we have pointed out, the reducing matrix is a special element of G (i.e., a Poincaré transformation) that parameterizes the gauge degrees of freedom other than

that of the Lorentz subgroup, i.e., the translations. Otherwise stated, the reducing matrix is a function of the coset parameters $\xi \in ISO(3,1)/SO(3,1)$. It is convenient to use a five dimensional matrix representation. We write the ten components of the Poincaré connection in the form

$$\Gamma_{\bar{B}}^{\bar{A}} = \begin{pmatrix} \Gamma_{\bar{b}}^{\bar{a}} & \Gamma^{\bar{a}} \\ 0 & 0 \end{pmatrix} \quad (11)$$

and a general Poincaré transformation as

$$P_{\bar{B}}^{\bar{A}} = \begin{pmatrix} \Lambda_{\bar{b}}^{\bar{a}} & \xi^{\bar{a}} \\ 0 & 1 \end{pmatrix}. \quad (12)$$

Clearly, the matrix that parameterizes the translations is given by

$$r_{\bar{B}}^{\bar{A}} = \begin{pmatrix} \delta_{\bar{b}}^{\bar{a}} & \xi^{\bar{a}} \\ 0 & 1 \end{pmatrix}. \quad (13)$$

We denote by $\bar{A} = (\bar{a}, \bar{5})$ the Poincaré group indices (instead of the greek ones, used above), in order to prevent confusion with the later on introduced $GL(R^4)$ indices, and by $A = (a, 5)$ the reduced (Lorentz) indices. Straightforward application of (8) gives for the reduced connection

$$\Gamma_B^A = \begin{pmatrix} \Gamma_b^a & e^a \\ 0 & 0 \end{pmatrix}, \quad (14)$$

where

$$\Gamma_b^a = \delta_a^{\bar{a}} \delta_b^{\bar{b}} \Gamma_{\bar{b}}^{\bar{a}} \quad \text{and} \quad e^a = \delta_a^{\bar{a}} [\Gamma^{\bar{a}} + \Gamma_{\bar{b}}^{\bar{a}} \xi^{\bar{b}} + d\xi^{\bar{a}}]. \quad (15)$$

This is the well know expression for the Lorentz connection and the tetrad field (see [17] or [18]). It remains to show the transformation properties of the new fields and of the coset parameters ξ^a . This can be done using equation (3). In matrix notation, the short form $\sigma' = \sigma(\xi')$ is to be interpreted as the fact that the transformed $r_{\bar{B}}^{\bar{A}}$ (seen as element of the Poincaré group) is again a pure translation. Using the transformation law in the form (9), i.e., explicitly

$$\tilde{r}_D^{\bar{C}} = \begin{pmatrix} \Lambda_{\bar{a}}^{\bar{c}} & a^{\bar{c}} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \delta_{\bar{b}}^{\bar{a}} & \xi^{\bar{a}} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} (L^{-1})^b_d & 0 \\ 0 & 1 \end{pmatrix}, \quad (16)$$

where the first matrix at the r.h.s. is a Poincaré transformation, and the third one a Lorentz transformation, a priori unrelated to each other, and then requiring that $\tilde{r}_D^{\bar{C}}$ is again a pure translation, i.e., of the form

$$\tilde{r}_D^{\bar{C}} = \begin{pmatrix} \delta_d^{\bar{c}} & \tilde{\xi}^{\bar{c}} \\ 0 & 1 \end{pmatrix}, \quad (17)$$

we find the condition $\Lambda^{\bar{a}}_{\bar{b}} = \delta^{\bar{a}}_{\bar{a}} \delta^{\bar{b}}_{\bar{b}} L^a_b$ (shortly $\Lambda = L$) and the transformation law

$$\tilde{\xi}^a = \Lambda^a_b \xi^b + a^a. \quad (18)$$

It is now easy to show that Γ^a_b and e^a in (15) transform indeed as Lorentz connection and Lorentz vector respectively.

In our special example with $ISO(3,1)$ as symmetry group, there is actually nothing nonlinear in the realization. Indeed, the relation $\Lambda = L$ allows us to identify the barred indices with the unbarred ones in the four dimensional subspace, as we have done in (18). In more complex examples, like the de Sitter [11] or the general linear [19] group, the transformation law (3) leads to a nonlinear dependence of the residual Lorentz transformation on the Lorentz part of the transformation of the initial symmetry group. Also, the reduced connection will depend nonlinearly on the coset parameters. We will see those features in the next sections, in the context of spontaneously broken gauge theories.

3 The de Sitter group

In this section, we present the gravitational gauge theory of the de Sitter group, spontaneously broken down to the Lorentz group, as presented by Stelle and West [11]. We do this not only because [11] seems to be the only article where a (conventional) Higgs sector is actually explicitly written down, but also because the de Sitter case, although nontrivial, is nevertheless nearer to conventional broken gauge theories, like the electroweak model or the nonlinear sigma-model and therefore presents a good starting point for the consideration of other symmetry groups which may present some particularities.

We start with a de Sitter (i.e., $SO(4,1)$) connection Γ^A_B (antisymmetric when we raise one index with the de Sitter metric $\eta_{AB} = \text{diag}(1, -1, -1, -1, -1)$) and construct the free Lagrangian density (or four form) $\mathcal{L}_0$ depending, ideally, on Γ^A_B only. (We will later see that the free Lagrangian can also depend on the Higgs field, in contrast to conventional Yang-Mills theories.) Then, in order to break down the symmetry, we have to introduce a Higgs field y^A carrying a de Sitter representation (in our case, y^A is simply a de Sitter vector). Note that the y^A are scalars under spacetime transformations. The Higgs potential $V(y^A)$ is chosen in a way that the groundstate $y^A(0)$ is invariant only under the Lorentz subgroup.

The first step is to find a decent (de Sitter invariant) characterization of such a groundstate. This is not very hard in this case, we may take

$$y^A(0)y_A(0) = -v^2, \quad (19)$$

where v^2 is a constant. Let us suppose that we have a Higgs sector that leads to (19) for the groundstate. (We will first consider the symmetry breaking and construct the Lagrangian afterwards.) The next step is to choose, from all the possible groundstates (19), the Lorentz invariant one

$$y^A(0) = (0, 0, 0, 0, v) \quad (20)$$

and to expand y^A in terms of new fields (which vanish in the groundstate) around $y^A(0)$. A convenient parameterization is given by

$$y^A = \begin{pmatrix} (\xi^a/\xi)(v + \varphi) \sin(\xi/v) \\ -(v + \varphi) \cos(\xi/v) \end{pmatrix}, \quad (21)$$

where $\xi = \sqrt{-\eta_{ab}\xi^a\xi^b} = \sqrt{-\xi_a\xi^a}$. Note that $y^A y_A = -(v + \varphi)^2$, i.e., φ gives us a measure of how far we are from the groundstate. It is the residual Higgs field of the theory. (We call y^A the (unbroken) Higgs field and φ the residual Higgs field, but occasionally refer to either of them as Higgs field.) Clearly, the remaining fields ξ^a are pure gauge, i.e., they can be transformed away. Indeed, performing a de Sitter gauge transformation with

$$\Lambda^A_B = \begin{pmatrix} \delta^a_b + (\cos(\xi/v) + 1)(\xi^a\xi_b/\xi^2) & -(\xi^a/\xi) \sin(\xi/v) \\ -(\xi_b/\xi) \sin(\xi/v) & -\cos(\xi/v) \end{pmatrix}, \quad (22)$$

the Higgs field reduces to the form

$$\tilde{y}^A = (0, 0, 0, 0, v + \varphi). \quad (23)$$

In order for our action to remain unaffected, we have to carry out the gauge transformation on all the fields, especially on the connection (we rescale the pseudo-translational part for dimensional reasons and take $e^a = v\Gamma^a_5$, since we intend to interpret e^a as tetrad field)

$$\Gamma^A_B = \begin{pmatrix} \Gamma^a_b & \frac{1}{v}e^a \\ \frac{1}{v}e_b & 0 \end{pmatrix}, \quad (24)$$

which under the transformation

$$\tilde{\Gamma}^A_B = \Lambda^A_C \Gamma^C_D \Lambda^{-1D}_B + \Lambda^A_C d\Lambda^{-1C}_B, \quad (25)$$

takes the form

$$\tilde{\Gamma}^A_B = \begin{pmatrix} \tilde{\Gamma}^a_b & \frac{1}{v}\tilde{e}^a \\ \frac{1}{v}\tilde{e}_b & 0 \end{pmatrix}, \quad (26)$$

where

$$\begin{aligned} \tilde{\Gamma}^a_b &= \Gamma^a_b + \frac{\cos(\xi/v) + 1}{\xi^2} (\Gamma^c_b \xi^a \xi_c + \Gamma^a_c \xi^c \xi_b) - \frac{\sin(\xi/v)}{v\xi} (\xi^a e_b - e^a \xi_b) \\ &\quad + \frac{\cos(\xi/v) + 1}{\xi^2} (\xi_b d\xi^a - \xi^a d\xi_b), \end{aligned} \quad (27)$$

$$\begin{aligned} \tilde{e}^a &= -e^a \cos(\xi/v) + v\Gamma^a_c \frac{\xi^c}{\xi} \sin(\xi/v) - \frac{1 + \cos(\xi/v)}{\xi^2} \xi^a e_c \xi^c \\ &\quad + \frac{1 + \sin \xi/v}{\xi^3} \xi^c d\xi_c \xi^a + \frac{\sin(\xi/v)}{\xi} d\xi^a. \end{aligned} \quad (28)$$

This result coincides with the one given in [11], after the replacement $\cos \rightarrow \cosh$ and $\sin \rightarrow \sinh$ and if one takes into account the difference in the sign convention. In [11], the signature is taken to be $\eta_{AB} = \text{diag}(-1, +1, +1, +1, -1)$, i.e., the symmetry group is $SO(3, 2)$, the anti-de Sitter group, and the global sign convention is opposite to ours.

Note that this result was achieved without the use of the cumbersome exponential parameterization that is used almost throughout the literature, including [11], and thereby, the painful manipulations associated with the use of the Campell-Hausdorff formula could be avoided.

If we compare with the last section, we recognize in (22) the reducing matrix, that is used to produce the nonlinear Lorentz connection (27) and the tensor (or rather vector) part (28). We also see that the coset parameters ξ^a appear now as the Nambu-Goldstone bosons of the theory. They are removed (gauged away) from the Higgs field and absorbed by the connection.

The remaining step is the construction of the Lagrangian. The appropriate Higgs sector has been constructed by Stelle and West. Introduce the following tensor:

$$g_{ik} = \frac{v^2}{y^C y_C} \left[\frac{1}{y^D y_D} (y^A D_i y_A)(y^B D_k y_B) - D_i y^A D_k y_A \right], \quad (29)$$

where $D_i y^A = y^A_{,i} + \Gamma^A_{Bi} y^B$. Clearly, g_{ik} is de Sitter gauge invariant. In the gauge (23), it takes the simple form

$$g_{ik} = \tilde{e}^a_i \tilde{e}^b_k \eta_{ab}. \quad (30)$$

We can now construct the following Higgs Lagrangian density:

$$\mathcal{L}_{higgs} = \sqrt{-g} \left[-\frac{1}{2y^C y_C} (y^B D_i y_B) (y^C D_k y_C) g^{ik} - V(y) \right], \quad (31)$$

with the Higgs potential

$$V = \frac{m}{2} y^A y_A + \frac{\lambda}{4} (y^A y_A)^2, \quad (32)$$

with m and λ positive constants. Obviously, the groundstate is characterized by the conditions

$$y_A y^A = -\frac{m}{\lambda} = -v^2, \quad (33)$$

and

$$y^A D_i y_A = 0. \quad (34)$$

Note that this last condition somewhat differs from the usual case. One would have expected a kinetic term of the form $(D_i y^A)^2$ and a groundstate condition $D_i y^A = 0$. This however would destroy the possibility of interpreting $\tilde{e}_m^a$ as tetrad field (and g_{ik} as metric), because then, in the groundstate the tetrad would vanish and certainly not be invertible.

In the gauge (23), the Higgs Lagrangian reduces to

$$\mathcal{L}_{higgs} = \sqrt{-g} \left[\frac{1}{2} \varphi_{,i} \varphi^{,i} + \frac{m}{2} (v + \varphi)^2 - \frac{\lambda}{4} (v + \varphi)^4 \right], \quad (35)$$

which is of the conventional form.

It remains to complete the theory with the Lagrangian for the gravitational field itself. In conventional theories, the Higgs field does not appear in the Yang-Mills sector. In gravity, this is not always possible, as we will see, because it is rather difficult to construct a Lagrangian four form using exclusively the gauge potentials. However, in the de Sitter case, Stelle and West have presented such a Lagrangian. It is of the form

$$\mathcal{L}_0 = \sqrt{-(F^{AB} \wedge F^{CD} \varepsilon_{ABCDE}) (F^{FG} \wedge F^{HI} \varepsilon_{FGHIJ}) \eta^{EI}}, \quad (36)$$

where F^{AB} is the curvature tensor of the de Sitter group which has the following components

$$F^{ab} = R^{ab} + \frac{1}{v^2} (e^a \wedge e^b), \quad F^{a5} = \frac{1}{v} (de^a + \Gamma^a_b \wedge e^b). \quad (37)$$

Going over to the gauge (23) (which means simply replacing Γ, e by $\tilde{\Gamma}, \tilde{e}$), where $\tilde{\Gamma}^a_b$ and $\tilde{e}^a$ are now the Lorentz connection and the tetrad, we recognize (dropping the $\sim$ for convenience) in R^{ab} the Lorentz curvature and in F^{a5} the torsion 2-form T^a .

There is one case where we can simplify (36). If the torsion vanishes, we simply get, omitting in the second line the Gauss-Bonnet divergence term

$$\begin{aligned}
\mathcal{L}_0 &= -F^{ab} \wedge F^{cd} \varepsilon_{abcd} \\
&= -2R^{ab} \wedge e^c \wedge e^d \varepsilon_{abcd} - e^a \wedge e^b \wedge e^c \wedge e^d \varepsilon_{abcd} \\
&= e \frac{8}{v^2} (R - \frac{3}{v^2}),
\end{aligned} \tag{38}$$

which is general relativity (in the first order formalism) with a cosmological constant. Unless some matter field is coupled to the Higgs field φ and to the Lorentz connection (as is the case with fermion fields), the theory remains in its groundstate $\varphi = 0$ and $T^a = 0$. The general case with non-vanishing torsion is difficult to handle with the non polynomial Lagrangian.

An alternative Lagrangian has also been proposed by Stelle and West, namely

$$\mathcal{L}_0 = F^{AB} \wedge F^{CD} \varepsilon_{ABCDE} y^E / v. \tag{39}$$

As before, using (37), you will find (in the gauge (23)), a curvature scalar term, a Gauss-Bonnet term and a cosmological constant term, each of them appearing, however, with a factor $(v + \varphi)$. Rescaling the scalar field by $1/v$ to make it dimensionless, and shifting it by one (to get a groundstate $\varphi = 1$ instead of $\varphi = 0$), the theory reveals itself to be very similar to Brans-Dicke theory (in a Riemann-Cartan framework). Note however that we cannot omit the Gauss-Bonnet term anymore. This term, of second order in the curvature, will eventually lead to dynamical torsion fields (in addition to those induced by the scalar field, see [11]) in the presence of spinning matter fields.

4 The Poincare group

The main difference between the de Sitter and the Poincaré group is the fact that the latter is not semi-simple. As a result, there is no Cartan metric available. This makes it rather difficult to construct invariants.

Let us begin by the introduction of a Higgs field y^a , carrying a (vector) representation of the Poincaré group, i.e., transforming as

$$y^a \rightarrow \Lambda^a_b y^b + a^a \tag{40}$$

under a Poincaré transformation (Λ^a_b, a^a) . For convenience, we do not distinguish notationally between a Lorentz and a Poincaré index (as we did in section 2).

The next step is to find a groundstate $y^a(0)$ that is Lorentz invariant, but breaks the Poincaré invariance. For the groundstate to be Lorentz invariant, it can only be characterized by a Lorentz invariant relation $\eta_{ab} y^a(0)y^b(0) = v^2$. Clearly, the only case where we have a Lorentz invariant solution to this is for $v^2 = 0$. It is the state

$$y^a(0) = 0. \quad (41)$$

This is indeed Lorentz invariant and breaks the Poincaré invariance.

To hold track with the de Sitter case, equation (41) corresponds to equation (20). We have now to parameterize a general state in terms of the groundstate, and write

$$y^a = \xi^a, \quad (42)$$

which is the counterpart of equation (21). This looks quite trivial, but it is the most general parameterization there is. Not surprisingly, it is pure gauge (as is any Poincaré vector). Otherwise stated, y^a is expressed entirely in terms of the Nambu-Goldstone bosons. There is no residual Higgs field in the theory. The transformation corresponding to the pseudo-translation (22) is now a pure translation (it is the $r^A_{\bar{B}}$ of section 2). Transforming the Higgs field and the connection, we find

$$\tilde{y}^a = 0, \quad \tilde{\Gamma}^a_b = \Gamma^a_b, \quad \tilde{\Gamma}^a \equiv e^a = \Gamma^a + d\xi^a + \Gamma^a_b \xi^b. \quad (43)$$

This is to be compared with (27) and (28).

We go on constructing the Higgs Lagrangian. What kind of a Higgs potential allows for a groundstate $y^a(0) = 0$? Well, the most simple solution is certainly $V(y) = 0$. This is not only short and elegant, but seems to be also the unique possible choice. As mentioned above, in Poincaré gauge theory, there are rather few objects to construct invariants. The Minkowski metric η_{ab} , although not related to a Cartan metric, nevertheless constitutes a natural ingredient of the theory, since it appears explicitly in the structure constants of the group algebra. It is therefore not unnatural to simply declare it as *invariant* under Poincaré transformations. You can also see it simply as a constant matrix. Similarly, you may see ε_{abcd} as Poincaré invariant.

With these objects, however, and the Poincaré vector y^a , the only scalar density you can construct is given by

$$\sqrt{-g} V(y) = \sqrt{-g} \lambda, \quad (44)$$

with a constant λ and with

$$g_{ik} = (D_i^{PG} y^a)(D_k^{PG} y^b) \eta_{ab}, \quad (45)$$

where D_i^{PG} denotes the Poincaré covariant derivative, $D_i^{PG} y^a = y_{,i}^a + \Gamma_{bi}^a y^b + \Gamma_i^a$. Note that this is not really a covariant derivative in the sense that it transforms in the same way as y^a . It has however the nice property that its vanishing in one gauge assures the vanishing in any gauge. In other words, it is a Lorentz vector. The metric g_{ik} is of course gauge invariant.

In (44), the potential V actually consists of an (almost) trivial constant λ . The only reason for which it is not trivial are the fields that appear inside the metric g_{ik} in the factor $\sqrt{-g}$. But in view of this, and considering (45), we should rather consider (44) as part of the kinetic term of the Higgs Lagrangian. We therefore conclude that $V(y) = 0$.

As to the kinetic part, we already found one candidate with (44), namely $\sqrt{-g}\lambda$. Other ideas, like

$$\varepsilon_{abcd} D^{PG} y^a \wedge D^{PG} y^b \wedge D^{PG} y^c \wedge D^{PG} y^d \text{ or } \sqrt{-g} g^{ik} D_i^{PG} y^a D_k^{PG} y_a$$

are only different forms or the same expression. No other expression containing only first derivatives of y^a can be formed with the ingredients we have at our disposal. Indeed, the next most simple candidate is the Riemannian curvature scalar for the metric g_{ik} . This contains higher derivatives of y^a (in the metric derivatives) and moreover, the Riemannian curvature can also be formed as a part of $\mathcal{L}_0$, as we know from traditional Poincaré gauge theory. It is then written as the sum of the teleparallel Lagrangian and the (Lorentz) curvature scalar.

The total, most general Higgs Lagrangian therefore has the form

$$\mathcal{L}_{higgs} = \sqrt{-g} \frac{\lambda}{4} g^{ik} D_i^{PG} y^a D_k^{PG} y_a = \sqrt{-g} \lambda. \quad (46)$$

In its first form, this looks like a conventional kinetic term for the Higgs field y^a , but one should not forget that during the variation, the metric has to be taken into account too. In the gauge (43), we simply get (omitting the $\sim$)

$$\mathcal{L}_{higgs} = e \lambda, \quad (47)$$

where e denotes the determinant of e_m^a . Summarizing, the Higgs potential is zero and the kinetic Higgs sector corresponds to a cosmological constant. This explains why

apparently, in Poincaré gauge theory, nobody included a Higgs sector. It was already there, but we did not recognize it as such.

It remains to complete the theory by constructing the gravitational sector. The only covariant quantity we can construct out of the Poincaré connection (we use again a five dimensional representation in the form (11)) is the Yang-Mills tensor

$$F^A_B = \begin{pmatrix} R^a_b & \tau^a \\ 0 & 0 \end{pmatrix} \quad (48)$$

where R^a_b is the Lorentz curvature and $\tau^a = d\Gamma^a + \Gamma^a_b \wedge \Gamma^b$. You can also check that the ε_{ABCDE} is again a tensor density under (12). With these objects alone, it is quite difficult, although not impossible, to construct Lagrangians. To be complete, we give an example:

$$\mathcal{L}_0 = \sqrt{(F^A_E \wedge F^B_F)(F^C_G \wedge F^D_H)\varepsilon_{ABCDI}\varepsilon^{EFGHI}}. \quad (49)$$

We do not know if it is possible to reduce this expression to something more easy to handle, or even to something useful, but it shows that there are, at least in principle, actions that are invariant under the Poincaré group, without the help of additional structures.

You may also look for actions formed out of the four dimensional parts R^a_b and τ^a . This has the advantage that, in addition to ε_{abcd} , you also have η_{ab} as invariant. However, no obvious scalar can be constructed with τ^a , due to its non homogeneous behavior under translations, and quantities constructed with the curvature, like $R^{ab} \wedge R_{ab}$ or $R^{ab} \wedge R^{cd}\varepsilon_{abcd}$ turn out to be total derivatives. (Of course, there is also a solution to that: Take the square of the first, add the square of the second and take the square root. The result is certainly not a total derivative.)

In order to construct more conventional, polynomial, Lagrangians, we see that the only way is to use the Higgs field also in the gravitational part of the Lagrangian. The Einstein-Cartan Lagrangian, for instance, is written as

$$\mathcal{L}_0 = D^{PG}y^a \wedge D^{PG}y^b \wedge R^{cd}\varepsilon_{abcd} \quad (50)$$

which is gauge invariant and reduces in the gauge (43) to

$$\mathcal{L}_0 = e^a \wedge e^b \wedge R^{cd}\varepsilon_{abcd}. \quad (51)$$

More generally, if we introduce the fields

$$E^a = D^{PG}y^a \quad \text{and} \quad T^a = dE^a + \Gamma^a_b \wedge E^b \quad (52)$$

then any Lagrangian constructed out of E^a , R^{ab} and T^a will be Poincaré invariant. These quantities are Lorentz tensors (i.e., invariant under translations). E^a plays the role of the tetrad field, as is clear in the gauge (43), where it reduces to $E^a = e^a$. In the same gauge, T^a reduces to the torsion tensor. In this way, you get the whole family of conventional Poincaré gauge theory Lagrangians.

More generally, one can argue that no Lagrangian can be constructed without the help of y^a that contains the translational gauge field Γ^a . You see this as follows: If the field y^a is contained in $\mathcal{L}_0$, then it appears necessarily only in the combination $D^{PG}y^a$. This however, in the specific gauge $y^a = 0$, reduces to Γ^a . The other way around, if the Lagrangian depends only on Γ^a_b and Γ^a , you can interpret Γ^a as $D^{PG}y^a$ in the gauge $y^a = 0$, since the Lagrangian is gauge invariant. This would mean that, in addition to its Poincaré gauge invariance, the Lagrangian would have to be invariant under the *replacement* $\Gamma^a \rightarrow D^{PG}y^a$, since else, it can not be gauge invariant with and without the y^a terms. Clearly, this is only possible when Γ^a is not contained at all in $\mathcal{L}_0$. Note that this is also the case with (49), which reduces immediately to its four dimensional parts, since $F^A_5 = 0$.

However, if $\mathcal{L}_0$ does not depend on Γ^a , i.e., on e^a in the final gauge, then we will not have an Einstein equation, which means that the energy-stress tensor of the matter fields will have to vanish. Such Lagrangians can therefore not be considered as physically acceptable. As a result, the Higgs field y^a is a necessary ingredient of Poincaré gauge theory, apart from its role in the symmetry breaking process.

Before we close this section, let us make some remarks about the groundstate of the theory. We have concluded earlier that the groundstate has to be described by $y^a = 0$, because it is the only choice that is Lorentz invariant. However, the groundstate can not be guessed simply from the form of the potential or from other arguments, it has to be the result of the field equations for the Higgs field. If you look at (46), you might get the impression that the groundstate is characterized by $D^{PG}_i y^a = 0$. This would be rather unfortunate, since this is just the quantity that is to be used as tetrad field. We should therefore check the situation more carefully.

Let us first consider the (conventional, but in gravity unlikely) case where the Lagrangian $\mathcal{L}_0$ does not depend on the Higgs field. Then, in the absence of matter field that couple to y^a , the field equation for y^a is easily derived by varying (46)

$$T^b_{ik} E^i_a E^k_b = 0, \quad (53)$$

or simply, transforming tangent space indices into spacetime indices using the tetrad,

$T^l_{ml} = 0$. The groundstate is thus characterized by equation (53), which is the analogue of equation (34) in the de Sitter case. This is clearly a Poincaré covariant relation (meaning that, if it holds in one gauge, then in any; not meaning that the left hand side is a Poincaré vector) and you may now go on and choose the state $y^a(0)$ in Lorentz invariant way.

In the usual case, the Lagrangian $\mathcal{L}_0$, and also any matter Lagrangian $\mathcal{L}_m$ will depend on y^a through the tetrad field $E^a = D^{PG}y^a$. In this case, it is easy to see that the equation arising from variation with respect to y^a is identically satisfied if the Einstein equation (i.e., the equation for E^a , or, equivalently, for Γ^a) is satisfied. Indeed, the y^a equation is just the covariant derivative of the E^a equation, see [17]. Therefore, no additional condition arises on the tetrad field that would exclude its invertibility.

In practice, the usual procedure is to first gauge the Nambu-Goldstone bosons away and to vary the action afterwards. Then, since y^a was pure gauge, the Higgs field does not appear anymore, and the only thing that reminds us of it is the cosmological constant (i.e., a mass term for the field e^a , the field corresponding to the broken gauge symmetry). We see that the fact that y^a is pure gauge is directly related to the fact that its field equation is identically satisfied and that after the symmetry breaking, no Higgs field remains in the theory.

There is only one question left: Can we choose $\lambda = 0$? Well, from a practical point of view, there is no problem with this. Theories without cosmological constant are quite commonly accepted. Choosing $\lambda = 0$ erases the last trace of the Higgs field and leaves y^a without kinetic term. In the spirit of conventional theories with dynamical symmetry breaking, even though y^a is of pure gauge nature, and we therefore can live without this term, one should not omit it. As kinetic term for the field y^a (look at it in the first form of equation (46)), one might even argue that the *natural* choice would correspond to $\lambda > 0$, i.e., to a positive cosmological constant.

5 The affine group

Considering the affine group as symmetry group of the gravitational interaction leads to the so called metric affine theory. We refer to [20] for a detailed presentation of its features. The nonlinear realization with the Lorentz group as stability subgroup, as well as references to earlier attempts can be found in [19].

To the affine group correspond 20 gauge fields Γ^α_β and Γ^α , $\alpha, \beta = 0, 1, 2, 3$. With the translational part Γ^α , one deals exactly as in the Poincaré case, i.e., introduces a Higgs field y^α , transforming as affine vector, then defines $E^\alpha = \Gamma^\alpha + dy^\alpha + \Gamma^\alpha_\beta y^\beta$ and so on. After gauging away the Nambu-Goldstone bosons, we have $E^\alpha = \Gamma^\alpha \equiv e^\alpha$. We are then left with the 20 fields Γ^α_β and e^α , a $GL(R^4)$ -connection and a $GL(R^4)$ -vector respectively.

In this section, we concentrate on the reduction of $GL(R^4)$ (shortly GL) to $O(3, 1)$. Let us briefly review the nonlinear realization process. As in section 2, we introduce a reducing matrix r^α_b with a mixed transformation behavior given by equation (9), where H^α_b is now a Lorentz and G^α_β a GL transformation. The Lorentz connection is then given by (8).

What is a suitable reducing matrix? Recall that r^α_a is itself an element of GL . In the Poincaré case, the reducing matrix was a pure translation. Similarly, in this case, it will be a general linear transformation that does *not contain* a Lorentz rotation. We know that infinitesimal Lorentz transformations are antisymmetric. Antisymmetric with respect to the Lorentz metric, more precisely. Because, if we take any (infinitesimal) general linear transformation ε^α_β , and look at a general metric tensor $g^{\alpha\beta}$, then this metric will be invariant if we have $\varepsilon^\alpha_\gamma g^{\gamma\beta} = -\varepsilon^\beta_\gamma g^{\alpha\gamma}$, i.e., if $\varepsilon^{\alpha\beta}$ is antisymmetric where the second index has been raised with $g^{\alpha\beta}$. It is a Lorentz transformation, by definition, only in the case where the invariant metric is the Minkowski metric. In that case, we say that ε^α_β is antisymmetric with respect to the Minkowski (or Lorentz) metric. In [7], the expressions pseudo-symmetric and pseudo-antisymmetric are used in this sense.

The infinitesimal Lorentz transformations being antisymmetric in the above sense, the infinitesimal form of our reducing matrices have to be symmetric. Exponentiating a symmetric matrix leads to a symmetric matrix. Therefore, we finally have the following characterization:

$$r^\alpha_a \eta^{a\beta} = r^\beta_a \eta^{a\alpha}. \quad (54)$$

This reduces the independent degrees of freedom contained in r^α_a to 10, as many as the parameterized gauge degrees of freedom.

The appearance of the Minkowski metric is unavoidable at this stage, it is the result of having chosen a specific stability subgroup. If we take $diag(1, 1, 1, 1)$ instead of η_{ab} , we will end up with an $O(4)$ symmetry. Unfortunately, with the exception of [7], little attention has been paid to this crucial point. One usually splits the GL generators G^α_β

into a *Lorentz* and a *symmetric* part $L^\alpha_\beta + S^\alpha_\beta$, without mentioning anything about a metric. The truth is, however, that you cannot split in this way the generators *before* the introduction of the Minkowski metric. This can already be seen from the concrete representation with the *GL*-generators taken in the form $x^k \partial_i$. You cannot extract the angular momentum generators from this, without the use of a metric. The geometric reason for this is the fact that in dimensions higher than three, rotations take place not *around an axis*, but *in a surface*, and therefore you have to know the geometry of that surface.

The requirement, that after a transformation (9), the reducing matrix remains symmetric in the sense of (54) leads to a nonlinear relation between the Lorentz part of the *GL* transformation G^α_β and the Lorentz transformation H^a_b in (9) (see [19]).

You can also define a metric through $g_{\alpha\beta} = r^a_\alpha r^b_\beta \eta_{ab}$, transforming as tensor under the *GL* group, but this seems rather unnatural at this point since our goal is just the contrary, i.e., to reduce *GL* quantities to Lorentz quantities.

We move on to the construction of a theory with dynamical breakdown of the symmetry group *GL* to the Lorentz stability subgroup $O(3,1)$.

Together with the fields e^α and Γ^α_β (recall that we consider the translational gauge symmetry to be already broken), we introduce 10 Higgs fields (as many as the gauge degrees of freedom we want to break) carrying a *GL* representation, in the form of a symmetric tensor $g_{\alpha\beta}$. We can, at this stage, also introduce a spacetime metric through

$$g_{ik} = e^\alpha_i e^\beta_k g_{\alpha\beta}. \quad (55)$$

We now look for a Higgs potential that is supposed to lead to a Lorentz invariant groundstate, but breaks the *GL* symmetry. The following considerations will help us to construct it. Recall the fact that *GL*, as opposed to the Lorentz group, is a very strong group, in the sense that we can diagonalize any symmetric matrix. More precisely, we can bring $g_{\alpha\beta}$ into one of the four forms $diag(\pm 1, \pm 1, -1, -1)$, depending on the signature of $g_{\alpha\beta}$ (see [20]). Note that we are not interested in the global sign of the metric, which is physically not relevant (different conventions are used in the literature with equivalent results). Therefore, if the groundstate turns out to have a specific signature, then the stability subgroup is fixed to the corresponding rotation group $O(3,1)$, $O(2,2)$ or $O(4)$. It seems quite impossible to write down a Higgs potential depending on the signature (expressed in terms of the $g_{\alpha\beta}$ components), but fortunately, there is one case where it is enough to know the sign of the determinant

in order to conclude for the signature. This is the case where the first two entries in $diag(\pm 1, \pm 1, -1, -1)$ are of opposite sign. Only then will the determinant be negative. The determinant is not an invariant, but it does not change sign under a GL transformation. This is the clue to the construction of the Higgs potential. All we have to do is to construct a Lagrangian that leads to a groundstate characterization $\det g_{\alpha\beta} < 0$.

This, however, turns out to be more difficult than it seems. The most simple scalar density that is invariant under GL and could eventually serve our purpose is the following

$$\sqrt{|g_{ik}|} V(g_{\alpha\beta}) = |e| \sqrt{|\det g_{\alpha\beta}| + \det g_{\alpha\beta}}.$$

Indeed, variation with respect to $g_{\alpha\beta}$ leads to

$$\delta(\sqrt{|g_{ik}|} V(g_{\alpha\beta})) = -|e| \sqrt{|\det g_{\alpha\beta}| + \det g_{\alpha\beta}} g_{\alpha\beta} \delta g^{\alpha\beta}, \quad (56)$$

and consequently, the groundstate is seemingly characterized by the condition

$$\det g_{\alpha\beta} < 0.$$

One could complete the Higgs sector in the following way

$$\mathcal{L}_{higgs} = \sqrt{|g_{ik}|} [g^{ik} D_i g_{\alpha\beta} D_k g^{\alpha\beta} - V(g_{\alpha\beta})]$$

It seems as if, as long as the non-metricity $Dg_{\alpha\beta}$ is zero, the theory would remain in its groundstate, i.e., the metric signature would be Lorentzian. This, however, is an illusion.

The problem with (56) is, that we will never get this equation in an isolated form, even if $Dg_{\alpha\beta} = 0$. You can directly check that an equivalent equation will be obtained from the variation with respect to e^α . Moreover, it is a general result [20] that in metric affine theory, one of the two equations arising from the variation with respect to the tetrad and to the metric respectively, is redundant (under the assumption that the matter equations and the Γ^α_β equations hold). The reason for this is that you can always gauge the metric into a non-dynamical, constant matrix (not necessarily of Minkowskian signature), or alternatively, you can gauge e^α_i into δ^α_i . Thus, one of both fields can be seen as pure gauge.

After these considerations, it is clear that the expression (56) will arise in both the e^α equation (i.e., ultimately, in the Einstein equation) and in the $g_{\alpha\beta}$ equation. There

will be no independent equation for $g_{\alpha\beta}$. The same will be true for whatever Higgs Lagrangian $\mathcal{L}_{higgs}(g)$ one might come up with.

Before we continue, let us note that the above argument also leads to the conclusion, that no Lagrangian can be constructed for the free gravitational fields without the help of $g_{\alpha\beta}$, apart from those that do not contain e^α , i.e., those who require a vanishing of the stress-energy tensor of the matter fields. Because, if $\mathcal{L}_0$ does not depend on the metric, its e^α equation too has to be identically satisfied (with the help of the other equations).

On the other hand, if we start with the symmetry group $SL^\pm$ (the subgroup of GL with $|\det G| = 1$ for $G \in GL$), this problem should not occur, since we cannot, in general, gauge e_i^α into δ_i^α . Indeed, under $SL^\pm$, we have an additional invariant given by $\det g_{\alpha\beta}$, and we can construct the following Higgs potential

$$\sqrt{|\det g_{ik}|} V = \sqrt{|g_{ik}|} [(\det g_{\alpha\beta}) + \frac{1}{2}(\det g_{\alpha\beta})^2]. \quad (57)$$

The only part of the equation resulting from the variation with respect to $g_{\alpha\beta}$ that will be independent from the other equations (especially from the e^α equation), is the part coming from the $\det g_{\alpha\beta}$ variation inside the brackets. Therefore, we are let to

$$\det g_{\alpha\beta} = -1. \quad (58)$$

You can now add a dynamical sector for the Higgs field $g_{\alpha\beta}$. The groundstate condition (58) clearly breaks the symmetry from $SL^\pm$ down to $O(3, 1)$. As long as the system remains in its groundstate, the metric will be related by an $SL^\pm$ gauge transformation to the Minkowski metric. We see that the introduction of additional (mani)fields is really not needed at this point. As opposed to the presentation of Hehl et al. in [20], in our approach, the nontrivial step is not the reduction from $SL^\pm$ to $O(3, 1)$, but rather the reduction $GL \rightarrow SL^\pm$.

How can we generalize (57) to the case of the GL group? Well, we have already argued that we will never get an equation for $g_{\alpha\beta}$ that is independent of the rest of the equations, including the matter field equations. Therefore, the necessary step is to include a new matter field that serves the purpose to break the equivalence between the other two equations. In other words, if we introduce an additional Higgs field φ , then we have the choice to consider either the set $(\Gamma_\beta^\alpha, e^\alpha, \varphi, \psi)$, where ψ summarizes the other matter fields eventually present, or alternatively the set $(\Gamma_\beta^\alpha, e^\alpha, g_{\alpha\beta}, \psi)$ as independent variables. For this to work, it is clear that φ should be of pure gauge

nature, that is, it should be possible to gauge it into something non-dynamical. After doing this, the gauge is fixed and we cannot trivialize anymore the tetrad or the metric, which will both take an independent status. Or the other way around, fixing the gauge by choosing the tetrad (or the metric) to be non-dynamical, the field φ becomes an independent dynamical quantity.

Clearly, φ cannot be a GL invariant. The next most simple candidate is a scalar density. This is also suggested by equation (57). It can easily be generalized if we choose φ to transform in a way that $\varphi^2 \det g_{\alpha\beta}$ is an GL invariant. In other words, φ has to be a scalar density of weight $+1$ under GL (and an ordinary scalar under spacetime transformations). Such a *dilaton* field has also been used in [20] in a similar approach. Note, however, that in contrast to [20], where the dilaton transforms as density under *conformal* transformations, whereas it is a scalar under GL transformations, we do not consider conformal transformations here.

Finally, we are ready to propose the following Higgs Lagrangian:

$$\begin{aligned} \mathcal{L}_{higgs} = \sqrt{|g_{ik}|} & \left[\varphi^2 (\det g_{\alpha\beta}) + \frac{1}{2} \varphi^4 (\det g_{\alpha\beta})^2 + \frac{1}{2} (\det g_{\alpha\beta}) g^{ik} D_i \varphi D_k \varphi \right. \\ & \left. + g^{ik} D_i g_{\alpha\beta} D_k g^{\alpha\beta} \right]. \end{aligned} \quad (59)$$

The covariant derivative of φ is defined by

$$D\varphi = d\varphi + \Gamma_{\alpha}^{\alpha} \varphi. \quad (60)$$

You can check that $D\varphi$ is again a scalar density of weight one. The metric determinant plays the role of the metric in the space of scalar densities, i.e., from two scalar densities φ, ψ , you form the scalar $\varphi \cdot \psi = (\det g_{\alpha\beta}) \varphi \psi$. In this sense, the first two terms in (59) are of the usual form of a Higgs potential $\varphi^2 + \frac{1}{2} \varphi^4$ for the field φ . (We have not included possible coupling constants in (59), since we are only interested in the main features of the symmetry breaking and not in specific models.)

The field equation for φ leads to the groundstate condition

$$(g + g^2 \varphi^2) \varphi = 0, \quad (61)$$

where $g = \det g_{\alpha\beta}$, which, under the assumption $\varphi, g \neq 0$, finally leads to

$$\det g_{\alpha\beta} < 0. \quad (62)$$

This, as we have argued, leads uniquely to the Lorentz signature of the metric. The same signature, through (55), is passed on to the physical spacetime metric.

Before we comment on the above approach, let us complete our procedure in a way similar to the previous sections. The gravitational Lagrangian may be taken as

$$\mathcal{L}_0 = \sqrt{|\det g_{\alpha\beta}|} \frac{e}{|e|} e^\alpha \wedge e^\beta \wedge R^\gamma{}_\kappa \varepsilon_{\alpha\beta\gamma\delta} g^{\delta\kappa}. \quad (63)$$

This is the direct generalization of the Einstein-Hilbert action to the metric affine framework. More general candidates can be found in [20]. The factor $e/|e|$ is necessary in order to insure invariance under a GL transformation with negative determinant, although it does not really change the field equations. From the groundstates characterized by (61), we choose

$$|\varphi(0)| = 1, \quad g_{\alpha\beta}(0) = \eta_{\alpha\beta}. \quad (64)$$

Any other choice, with the exception of $\varphi = 0$ or $g = 0$ is related to this one by a GL transformation. The first equation in (64) reduces the symmetry from GL to $SL^\pm$, and the second one finally leaves us with $O(3,1)$ as residual symmetry group. As before, we will use latin indices again for the Lorentz quantities. The next step is the parameterization of a general state in terms of the Nambu-Goldstone bosons and the residual Higgs fields. We were not able to find a very elegant solution to this, but let us write down the following attempt:

$$g_{\alpha\beta} = r^a{}_\alpha r^b{}_\beta \tilde{\eta}_{ab}, \quad (65)$$

with

$$\tilde{\eta}_{ab} = \text{diag}(1 + \mu, -1 + \nu, -1, -1). \quad (66)$$

The Nambu-Goldstone bosons $r^a{}_\alpha$ correspond to the reducing matrix of the nonlinear realization approach and are, as such, elements of GL . They also contain the parameterization of the (pure gauge) field φ (in their determinant). Since only those metrics can be related by a GL transformation to the Minkowski metric η_{ab} , that possess the correct signature, in order to describe a general metric, we have to introduce two *signature* functions μ and ν . They are the residual Higgs fields and cannot be gauged away. Clearly, they vanish in the groundstate. The form (66) is a modification of a similar parameterization proposed in [19, 28, 29].

In practice, (66) is not really useful, in the sense that it does not allow us to express the Lagrangians (59) and (63), after gauging $r^a{}_\alpha$ away, in terms of μ and ν only, i.e., to separate the non-dynamical Minkowski metric from $\tilde{\eta}_{ab}$ and to carry out the variation with respect to μ and ν only. In view of this, nothing is really gained by the

gauge transformation and we can equally well use (59) and (63) in their initial form. Eventually, one has to come up with a better parameterization, that allows to express the metric directly in terms of the residual Higgs field, in a way that we explicitly see, for instance, the terms of the form $\mu_{,i}\mu^{,i}$ that are contained in the last term of (59).

This, in principal, completes the analysis of the symmetry breaking mechanism of metric affine gauge theory. You can now go on and consider concrete models. The main features will be that, as long as nothing couples to the symmetric part of the connection (i.e., if there are no matter fields that possess a so called hypermomentum, see [20]), the theory will remain in its groundstate and the signature will always be Lorentzian. Only fields with hypermomentum will provide the kinetic terms in (59) with a source and could be able to change the signature of $g_{\alpha\beta}$ and therefore, ultimately, of g_{ik} .

One can also consider the possibility of changing $\mathcal{L}_0$ by a factor, say, $(\det g_{\alpha\beta})\varphi^2$ (which is a scalar). This is actually preferable, since, then, you can express $\mathcal{L}_0$ without the help of the inverse metric $g^{\alpha\beta}$ and without dividing by $\det g_{\alpha\beta}$, making it suitable for the study of eventual signature changes. Similar as in the approach of Hehl et al., but with $g\varphi^2$ playing the role of the scalar field, one is led to Brans-Dicke type theories. The analysis of concrete models, however, is related to the description of matter fields with hypermomentum, which will in most cases include spinor representations of GL (manifolds), and is therefore beyond the scope of this article.

Let us also note that, in contrast to the approach in [20], the nonmetricity is not on the same level as the torsion or the curvature. It should appear explicitly only in the Higgs sector of the theory, as it does in the last term of (59). The Lagrangian $\mathcal{L}_0$ should, consequently, be constructed from the torsion and the curvature tensors, which are the Yang-Mills tensors corresponding to the original gauge fields of the affine group. This does not prevent the nonmetricity (or rather the symmetric part of the connection, to which the nonmetricity reduces when $g_{\alpha\beta} = \eta_{\alpha\beta}$) to appear, even as a dynamical field, implicitly in terms like $R^\alpha_{\alpha ik} R^\beta_{\beta}{}^{ik}$.

Some critical remarks are at order considering the approach presented in this section. The scalar density φ was originally included to break the symmetry of GL down to $SL^\pm$. This would be the case, if one were let to a nonzero groundstate described by

$$|\varphi(0)| = \alpha, \quad (67)$$

for some (nonzero) constant α . Then, clearly, the residual symmetry is $SL^\pm$. However, (59) does not really do this job. Instead, it leads to (61), which does not exclude the

value $\varphi = 0$. Only if we assume that $\varphi \neq 0$ (in other words, if we assume that GL is broken down to $SL^\pm$), we get the condition $g < 0$, which breaks the symmetry down to $O(3, 1)$. Thus, the role of φ is actually to allow us to write down the Lagrangian (57) in a GL invariant form and break the symmetry down to the Lorentz group, under the assumption that φ possesses a nonzero groundstate. It does not really determine that groundstate. The usual argument from $\varphi^2 + \varphi^4$ Higgs theories, namely that the value $\varphi = 0$ does not represent a minimum, but rather a maximum of the potential, and therefore is not a stable groundstate, is not valid in our case, because whether we are dealing with a maximum or a minimum depends directly on the sign of the metric determinant in (61), which is what we want to determine in the first place.

In the dilaton approach in [20], the situation is quite different, because the dilaton field is used not to determine the signature of the metric, but merely to break the (additional) conformal symmetry of the theory.

[There is actually a source of confusion in [20]: The expressions local scale transformations, or dilations, refer to GL transformations of the special form $G^\alpha_\beta = \Omega \delta^\alpha_\beta$. We have adopted this vocabulary, and consequently, call our field φ a *dilaton* (since the effect of a general G^α_β on φ is identical to the effect of a dilation with $\det G = \det \Omega$). In [20], however, the so-called dilaton field (section 6) is actually a GL -scalar (since it is invariant under the dilation $L = 0, F = C = -1$, in the notation of [20]), and the broken symmetry is the conformal symmetry, not the scale, or dilation, invariance. More serious than this linguistic problem is the fact that the authors, in the subsequent section (section 6.5), consider the dilational part of GL as already broken by the (so-called) dilaton, and concentrate on the breakdown of $SL^\pm$ (or $T \ltimes SL$ in their notation). This is obviously mistaken, because even if the (so-called) dilaton takes a nonzero groundstate value, the broken symmetry will be the conformal one, leaving us therefore with the full, unbroken GL symmetry group. Clearly, one cannot break the dilation invariance using a GL invariant field.]

This point needs further clarification. In contrast to the case of the translations, where the mere existence of a Poincaré (or affine) vector y^a necessarily leads to a symmetry breaking, independently of the choice of the groundstate, in the case of the dilaton, the symmetry is only broken if the groundstate is different from zero. Otherwise stated, any vector y^a is related by a translation to the groundstate $y^a = 0$, but only nonzero values for φ are related by a GL transformation to the groundstate value, say, $|\varphi| = 1$.

One will thus have to come up with a potential that assigns a nonzero groundstate value to φ . This can only be done if we have expressions where φ does not appear

in the combination $\varphi^2 \det g$, but this makes it rather difficult, if not impossible, to construct invariants. New ideas are needed here.

For the moment, we will present a solution to the above problem, which is based on the introduction of a scalar field ψ , in addition to the dilaton φ . Consider the following Higgs Lagrangian:

$$\mathcal{L}_{higgs} = \sqrt{|g_{ik}|} [\psi(g\varphi^2 + 1) + \psi(g^{ik} D_i g_{\alpha\beta} D_k g^{\alpha\beta})], \quad (68)$$

with $g = \det g_{\alpha\beta}$. The field equations for ψ and φ read

$$g\varphi^2 + 1 + D_i g_{\alpha\beta} D^i g^{\alpha\beta} = 0 \quad (69)$$

$$\varphi g \psi = 0. \quad (70)$$

If the nonmetricity vanishes (which will be the case, again, if no matter fields couple to the symmetric part of the connection, i.e., if the kinetic term for $g_{\alpha\beta}$ has no source term), the first equation leads to the required groundstate characterization

$$g\varphi^2 = -1. \quad (71)$$

This excludes explicitly the values $\varphi = 0$ and $g = 0$. Therefore, the symmetry, this time, has really been broken down to $O(3,1)$, without any ad hoc assumptions. Clearly, the metric determinant has to be negative, and therefore, the signature of $g_{\alpha\beta}$ is Minkowskian and we can choose the groundstate $g_{\alpha\beta}(0) = \eta_{\alpha\beta}$. Any other choice, solution to (71), would be related by a GL transformation to this one and therefore be equivalent. From (71), you then also conclude that $|\varphi| = 1$, and from (70) that $\psi = 0$. This last relation is rather interesting, since it means that in the groundstate, the whole Higgs sector is actually zero, and therefore does not contribute to the Einstein and Cartan equations. In other words, as long as the nonmetricity is zero, the theory will be exactly equivalent to Poincaré gauge theory. Especially, there will be no contribution to the stress-energy tensor from the Higgs field. This has also the consequence that, as opposed to usual spontaneously broken theories, the gauge fields corresponding to the broken group generators will not acquire mass, since the term that is responsible for this, the last term in (68), vanishes together with ψ .

In (68), no kinetic terms for φ or ψ have been included. This is also not necessary, since the actual Higgs field is $g_{\alpha\beta}$. The role of the field φ is merely to form a GL invariant expression with the metric determinant, and ψ is now the field that is introduced to break the equivalence of the tetrad and metric equations, i.e., to get an

independent equation for the symmetry breaking mechanism. You may also see ψ as a Lagrange multiplier.

The Lagrangian (68), although it may appear artificial, is nevertheless the first Lagrangian that *really* does, what it is supposed to do, namely to break down the general linear group to the Lorentz group. It remains to see, if simpler solutions exist to this problem, that allow us to possibly eliminate one of the additional fields φ or ψ .

On the other hand, one could also consider to start right from the beginning with a theory based on the special affine group and avoid the problems we just described. This, however, seems rather unnatural and also breaks the direct link that exists between GL theories and general relativity. Indeed, a strong argument in favor of GL is the fact that we can gauge the tetrad into its trivial form δ_i^α , and then *identify* the spacetime indices with the GL indices. In order to maintain the gauge in the form $e_i^\alpha = \delta_i^\alpha$, it is necessary, every time we change the coordinate system, to perform, at the same time, a GL transformation with matrix $\partial\tilde{x}^i/\partial x^k$. Under such a combined transformation, $\Gamma_{\beta i}^\alpha$ (then written as Γ_{li}^k) will transform as a general relativity connection. Note that this also shows that, in a certain way, the gauge approach presented in this article, contains as a subcase the spacetime based gauge approach of [22] mentioned in the introduction. Indeed, fixing the gauge in the above way leads to a residual invariance in the form of the transformations considered in [22], and moreover, tangent space G/H , through $e_i^\alpha = \delta_i^\alpha$, is identified with the spacetime manifold, an assumption that is made in [22] and similar theories (e.g., [13, 14]) right from the beginning.

Also, the Lagrangian (63) will, in this gauge, essentially reduce (up to torsion and nonmetricity parts contained in the connection) to the Einstein-Hilbert Lagrangian. This relation to general relativity is completely lost if we start with the special linear group. Note also, that theories based on the special affine group will lead to Brans-Dicke type theories, because we can gauge the metric $g_{\alpha\beta}$, even if it is in its groundstate signature, only up to a conformal factor into the Minkowski metric, leaving us therefore with a residual scalar Higgs field.

Finally, and most importantly, we remind that the whole procedure, as described in this section, only works for the Lorentz stability subgroup. It is not possible, in a similar manner, to write down a Lagrangian that leads to a residual $O(4)$ or $O(2, 2)$ symmetry, since in those cases, the mere knowledge of the sign of the metric determinant is not enough to determine its signature. In those cases, one would have to write down a Lagrangian in terms of a direct expression of the signature (which is a scalar), expressed in terms of the metric components. This may be possible, but it

is not obvious to us, how this can be done in an simple way. Generally, the signature can be expressed in terms of the four eigenvalues E_i of the metric, normalized to one, $e_i = E_i/|E_i|$. Those eigenvalues, however, are the solutions to a fourth order equation, and as such, highly nontrivial expressions in $g_{\alpha\beta}$.

It seems therefore at least unlikely that a reasonable Lagrangian can be constructed that assigns a groundstate signature other then the Lorentzian one to the spacetime metric.

Before we close this section, in order to avoid misunderstandings, let us clarify the critical remarks we made towards the references [12, 21] on one hand and [20] on the other hand. We do not claim that there is anything wrong with the symmetry breaking mechanism based on the introduction of manifields, as proposed in [12] and [21]. It can be seen as an alternative way to trigger the symmetry breakdown from SL down to the Lorentz group, and as such, should be completed by a suitable mechanism for the breakdown $GL \rightarrow SL$. What we claim is that, if one tries to perform the step $GL \rightarrow SL$ with the help of a dilaton field in the way presented in this section, then the symmetry will break down to the Lorentz group anyway, and this makes the manifold superfluous. Nevertheless, the symmetry breaking mechanism described in [12, 21] is interesting by itself, and moreover, the infinite dimensional representations of the special linear group described in those references will be needed anyway, if we want to complete our theory (in a GL -covariant way) with spinor fields. Thus, the manifields will enter, not as Higgs fields, but simply as matter fields, as generalization of the Dirac (and similar) Lagrangians. On the other hand, the (so-called) dilaton introduced in [20] serves a different purpose, namely to break the conformal invariance of the theory's groundstate. Also here, the approach presents its own interests, in the context of theories with asymptotical conformal invariance at high energies, as described in [20]. However, in addition to the conformal symmetry, we will also have to break the dilation invariance, i.e., include a mechanism for the $GL \rightarrow SL^\pm$ breakdown.

Summarizing, in none of the above references, the symmetry breaking mechanism for the reduction $GL \rightarrow O(3, 1)$ is complete. Moreover, our approach is preferable in the sense that it clarifies the role of the metric tensor, and assigns to it the interpretation suggested by the nonlinear realization approach in [19]. However, it remains to see if it is possible, either to avoid one of the additional fields, or, alternatively, to give a physical interpretation to those fields. As to the second possibility, one might consider φ and ψ as Higgs fields responsible for the breakdown of additional symmetries eventually present in $\mathcal{L}_0$. Possible candidates are, for instance, the projective symmetry,

or, in the manner of Hehl et al., conformal invariance.

In this article, we will consider yet another possibility, namely to enlarge the symmetry group to the special linear group $SL(R^5)$, in the hope to combine the (good) features of the de Sitter theory with the (seemingly incomplete) features of the affine theory.

6 The group $SL(R^5)$

In this section, G^A_B ($A, B = 0, 1, 2, 3, 5$) denotes the elements of $SL(R^5)$ and greek indices denote the four dimensional part of $SL(R^5)$ quantities, i.e., $A = (\alpha, 5)$ etc. The algebra of $SL(R^5)$ is given by the following relations:

$$[L^A_B, L^C_D] = i(\delta^A_D L^C_B - \delta^C_B L^A_D). \quad (72)$$

The trace of the generators vanishes, $L^A_A = 0$, as a result of $\det G^A_B = 1$. In order to get a clearer picture about the contents of $SL(R^5)$, let us introduce the de Sitter metric $\eta_{AB} = \text{diag}(1, -1, -1, -1, -1)$. Then, the commutation relations can be written in the form

$$\begin{aligned} [L_{\alpha\beta}, L_{\gamma\delta}] &= i(\eta_{\alpha\delta} L_{\gamma\beta} - \eta_{\gamma\beta} L_{\alpha\delta}), \\ [P_\alpha, L_{\gamma\delta}] &= i(\eta_{\alpha\delta} P_\gamma), \quad [K_\beta, L_{\gamma\delta}] = -i(\eta_{\gamma\beta} K_\delta), \\ [P_\alpha, D] &= -iP_\alpha, \quad [K_\alpha, D] = iK_\alpha, \\ [P_\alpha, K_\beta] &= -i(\eta_{\alpha\beta} D - L_{\alpha\beta}), \\ [P_\alpha, P_\beta] &= [K_\alpha, K_\beta] = [D, L_{\alpha\beta}] = 0, \end{aligned} \quad (73)$$

with $P_\alpha = L_{\alpha 5}$, $K_\alpha = L_{5\alpha}$, $D = -L_{55}$. Replacing the $GL(R^4)$ generators $L_{\alpha\beta}$ with the Lorentz generators $L_{[\alpha\beta]}$, (73) takes the form of the algebra of the conformal group $SO(4, 2)$. In view of this, we will refer to P_α, K_α as generators of the pseudo-translations G^α_5 , and the pseudo-inversions G^5_α (more precisely, pseudo-special conformal transformations), respectively.

The group $SL(R^5)$ has also been considered in [30] in the context of the nonlinear realization approach,

Instead of the de Sitter metric, we introduce now a general Higgs metric g_{AB} , an $SL(R^5)$ -tensor, which may be parameterized as follows

$$g_{AB} = \begin{pmatrix} g_{\alpha\beta} + \varphi^2 k_\alpha k_\beta & \varphi^2 k_\alpha \\ \varphi^2 k_\beta & \varphi^2 \end{pmatrix}. \quad (74)$$

The notation $g_{55} = \varphi^2$ is used to be consistent with the notation of the previous section, as will become clear later. Although it turns out that g_{55} is actually positive in the gauge we are interested in, and more generally, we are not interested in the global sign of the metric, for the moment, there is no restriction on the sign of φ^2 . From (74), we find

$$\det g_{AB} = \varphi^2 \det g_{\alpha\beta}, \quad (75)$$

and, if the determinant does not vanish, the inverse metric reads

$$g^{AB} = \begin{pmatrix} g^{\alpha\beta} & -k^\alpha \\ -k^\beta & k^2 + 1/\varphi^2 \end{pmatrix}, \quad (76)$$

where $g^{\alpha\beta}$ is the inverse of $g_{\alpha\beta}$ and $k^\alpha = g^{\alpha\beta}k_\beta$, $k^2 = k_\alpha k^\alpha$. In addition to g_{AB} , we introduce a further Higgs field, a $SL(R^5)$ -vector y^A , requiring the condition

$$g_{AB} y^A y^B \neq 0. \quad (77)$$

With 20 Higgs fields (g_{AB}, y^A) , underlying one constraint, it should be possible to reduce the gauge group from the 24-parameter group $SL(R^5)$ to the 6 parameter Lorentz group.

Similar to the de Sitter case (see equ. (29)), we introduce the spacetime metric

$$g_{ik} = -\frac{v^2}{y^2} \left[\frac{1}{y^2} (y_A D_i y^A)(y_B D_k y^B) - D_i y^A D_k y^B g_{AB} \right], \quad (78)$$

where, as opposed to the de Sitter case, special care has to be given to the index positions ($y_A D_i y^A \neq y^A D_i y_A$). Note also that, in view of (29), the condition (77) actually has also been assumed (quietly) by Stelle and West, excluding thereby, for instance, the value $\varphi = -v$ in (23).

Let us also parameterize the $SL(R^5)$ connection as

$$\Gamma^A_B = \begin{pmatrix} \Gamma^\alpha_\beta & \frac{1}{v} e^\alpha \\ B_\beta & A \end{pmatrix}. \quad (79)$$

The constant v is a length parameter that appears explicitly in the Higgs Lagrangian. Note that an $SL(R^5)$ invariant is given by the trace Γ^A_A .

We propose the following Lagrangian

$$\mathcal{L}_{higgs} = \sqrt{|g_{ik}|} [(y_A D_i y^A)(y_B D_k y^B) g^{ik} + D_i g^{AB} D_k g_{AB} g^{ik} - V(y, g)], \quad (80)$$

with the Higgs potential

$$V(y, g) = (1 + \det g_{AB})(g_{AB}y^A y^B - v^2) \quad (81)$$

Just as in the affine case, $V(y, g)$ is actually more an interaction of the different Higgs fields than a true potential. Recall that with the special linear group, the metric can be transformed into a non dynamical constant matrix, up to a conformal factor. Therefore, the only truly dynamical quantity in g_{AB} , leading to an independent equation, is $\det g$, which is invariant under $SL(R^5)$. Thus, (81) leads to the groundstate characterization

$$g_{AB}y^A y^B = v^2. \quad (82)$$

Before we analyze the symmetry breakdown in detail, we will try to make contact with the results of the previous section. In view of (82), we can choose the groundstate

$$y^A(0) = (0, 0, 0, 0, y^5(0)), \quad (83)$$

where the value $y^5(0)$ will depend on the groundstate value of g_{55} . Let us choose $y^5(0) = v$ and $g_{55}(0) = 1$. As in the de Sitter case, we can gauge a general state y^A into the form

$$y^A = (0, 0, 0, 0, y^5). \quad (84)$$

This leaves us with a residual gauge invariance

$$G^A_B = \begin{pmatrix} G^\alpha_\beta & 0 \\ q_\beta & s \end{pmatrix}, \quad (85)$$

with the only restriction $s \det G^\alpha_\beta = 1$. Thus, G^α_β is a $GL(R^4)$ transformation. Let us not worry about the pseudo-inversions q_α for the moment. Consider the transformation

$$G^A_B = \begin{pmatrix} \delta^\alpha_\beta & 0 \\ 0 & s \end{pmatrix}. \quad (86)$$

This is not a $SL(R^5)$ transformation and has therefore to be accompanied by a $GL(R^4)$ transformation with inverse determinant, but nevertheless, it is instructive to see the transformation properties of the various fields under (86). For the metric components (74) we find

$$\varphi^2 \rightarrow \varphi^2/s^2, \quad g_{\alpha\beta} \rightarrow g_{\alpha\beta}, \quad k_\alpha \rightarrow s k_\alpha, \quad (87)$$

and for the connection (79)

$$\Gamma^\alpha_\beta \rightarrow \Gamma^\alpha_\beta, \quad e^\alpha \rightarrow s^{-1} e^\alpha, \quad B_\alpha \rightarrow s B_\alpha, \quad A \rightarrow A - s^{-1} ds. \quad (88)$$

Note also that g_{ik} is an invariant under (86). This is a consequence of the fact that g_{ik} is actually not only $SL(R^5)$, but also $GL(R^5)$ invariant. Let us recall the general classification of conformal transformations (in the metric affine framework) by Hehl et al. [20]. Consider a function Ω and constants C, L, F . We call a *projective* transformation the following

$$\Gamma^\alpha_\beta \rightarrow \Gamma^\alpha_\beta - C \delta^\alpha_\beta d(\ln \Omega). \quad (89)$$

By (*pure*) *conformal* transformation, we mean the following

$$g_{\alpha\beta} \rightarrow \Omega^L g_{\alpha\beta}, \quad (90)$$

and finally, a *dilation* or *scale transformation* refers to a $GL(R^4)$ gauge transformation with

$$G^\alpha_\beta = \Omega^F \delta^\alpha_\beta. \quad (91)$$

Under the combined transformation (89)-(91), we find

$$\Gamma^\alpha_\beta \rightarrow \Gamma^\alpha_\beta - (C + F) \delta^\alpha_\beta d(\ln \Omega), \quad g_{\alpha\beta} \rightarrow \Omega^{L-2F} g_{\alpha\beta}, \quad e^\alpha \rightarrow \Omega^F e^\alpha. \quad (92)$$

Note that we have slightly changed the scheme of [20], in order to get a clearer separation of the different parts. The constant C in [20] (which we will denote by C_{hehl}) is related to our C by $C_{hehl} = C + F$. The other constants are identical to those of Hehl et al.

On the occasion, let us note that the so-called dilaton in [20] transforms as $\sigma \rightarrow \Omega^{-L/2} \sigma$, i.e., it is invariant under dilations, as we have noticed in the previous section, and especially, it cannot be used to break the dilational invariance. On the other hand, our dilaton φ introduced in (59) transforms with the parameter F and is therefore a true dilaton.

Comparing (92) with (87) and (88), we see that the transformation (86) can be interpreted as a conformal transformation with parameters

$$F = -1, \quad C = 1, \quad L = -2. \quad (93)$$

Especially, we see that a dilaton φ is automatically present in the theory. Indeed, the Higgs potential (81), in the gauge (84), using (75) takes the form

$$V = \psi(1 + \varphi^2 \det g_{\alpha\beta}), \quad (94)$$

with $\psi = g_{AB} y^A y^B - v^2 = g_{55} y^5 y^5 - v^2$. In view of the groundstate $y^5(0) = v$, $g_{55}(0) = 1$, you can parameterize, for instance, a general state by $y^5 = (1/\sqrt{g_{55}}) \sqrt{v^2 + \psi}$,

which leads to (94) immediately. The important thing is that, independently of the parameterization we choose, the factor ψ is a scalar, whereas φ^2 is a dilaton. Thus, (94) is identical to the Higgs potential used in (68).

The group $SL(R^5)$ therefore provides a natural explanation for the appearance of the dilaton and the scalar field which had to be introduced by hand in the framework of the affine group. In contrast to (68), the Lagrangian (80) contains kinetic terms also for φ and ψ . This is quite natural, since now those fields appear as parts of the true Higgs fields g_{AB} and y^A , while in (68), they played only a secondary role.

Let us now return to (80) and (81) and complete the analysis of the symmetry breaking mechanism. As explained before, the only truly dynamical part of g_{AB} is contained in its determinant. Therefore, under the groundstate conditions

$$y_A D_i y^A = 0, \quad D_i g_{AB} = 0, \quad (95)$$

the first of which is the direct generalization of the de Sitter case (34), we find the following equations

$$g_{AB} y^A y^B = v^2, \quad y^A (\det g_{AB} + 1) = 0. \quad (96)$$

In view of the first equation, there will certainly be a non-zero component of y^A , and therefore we find $\det g_{AB} = -1$. As before, we choose the groundstate

$$y^A = (0, 0, 0, 0, v). \quad (97)$$

Then, the first equation in (96) leads to $g_{55}(0) = 1$. Therefore, from $\det g_{AB}(0) = -1$ and with the help of (75), we are led to $\det g_{\alpha\beta}(0) = -1$, which means that we may choose the groundstate $g_{\alpha\beta}(0) = \eta_{\alpha\beta}$, the Minkowski metric with Lorentz signature. This leaves us with

$$G^A_B = \begin{pmatrix} \Lambda^\alpha_\beta & 0 \\ q_\beta & 1 \end{pmatrix}, \quad (98)$$

as residual gauge invariance, with Λ a Lorentz transformation. It is easy to show that the remaining parts of the metric, $g_{\alpha 5}$, can be gauged away with a pseudo-inversion q_β . This leaves us with the groundstate

$$g_{AB}(0) = \text{diag}(1, -1, -1, -1, 1), \quad y^A(0) = (0, 0, 0, 0, v), \quad (99)$$

which is clearly Lorentz (and only Lorentz) invariant.

As was the case with $GL(R^4)$, the parameterization of a general state is not trivial. Let us concentrate on those parts that cannot be transformed away. In other words,

also in a general state, we can always fix the pseudo-translational and the pseudo-inversional gauge by requiring $y^\alpha = 0$ and $g_{\alpha 5} = 0$. Then, one might consider to use the G^5_5 gauge to fix $g_{55} = 1$. Alternatively, one might fix y^5 to v (not both, however). Let us take the first choice, and parameterize $y^5 = \sqrt{v^2 + \psi}$. At this point, the Higgs potential takes the form

$$V = \psi(1 + \det g_{\alpha\beta}), \quad (100)$$

and the residual invariance group is $SL(R^4)$. Note also that (78) takes the expected form $g_{ik} = e_i^\alpha e_k^\beta g_{\alpha\beta}$. The dilaton seems to have disappeared, but remember that we can transform $g_{\alpha\beta}$ only up to a conformal factor into a constant matrix. Thus, for the four dimensional part of the metric, using a similar parameterization as in (65), (66), and fixing the gauge by transforming away the Nambu-Goldstone bosons, we can write

$$g_{ab} = \varphi^2 \tilde{\eta}_{ab}, \quad (101)$$

with

$$\tilde{\eta}_{ab} = \text{diag}(1 + \mu, -1 + \nu, -1, -1). \quad (102)$$

We use latin indices again, to underline the residual Lorentz invariance. Again, we can consider φ^2 to be positive, since the global sign of the metric is not important.

Thus, the broken, Lorentz invariant Lagrangian contains 4 residual Higgs fields, φ, μ, ν and ψ . At this stage, they transform simply as scalar fields. Their kinetic terms are contained in (80), but again, it will be difficult to find an explicit form in terms of μ and ν and it might be preferable go one step back and use an $SL(R^4)$ invariant form.

The full analysis of the $SL(R^5)$ theory is beyond the scope of this article. The theory, compared to the metric affine theory, contains five additional fields, B_a and A which will have to be interpreted and whose coupling to matter fields will have to be discussed. The incorporation of spinor fields in our theory can be done using the SL spinor representations of Ne'eman and Sijacki [12, 21]. As opposed to the metric affine theory, where this is only possible after reducing $GL(R^4)$ down to $SL(R^4)$, a fully $SL(R^5)$ -invariant Lagrangian with spinor matter can thus be written down.

7 Nambu-Goldstone fermions and $GL(R^4)$ invariant Dirac Lagrangian

Finally, we will briefly sketch a quite different model of a dynamically broken gravitational theory. Although the theory, as it will be presented here, is still incomplete, especially with respect to the interpretation of the tetrad field and the incorporation of translations, it is nevertheless rather promising under several aspects. First, it enables us to write down a $GL(R^4)$ covariant equation for a simple vector field k^α (as opposed to an infinite dimensional $SL(R^4)$ spinor), that, after the symmetry breaking, reduces to the usual form of the Dirac equation without any ad hoc assumptions. Moreover, the fermion fields appear under a very different and rather instructive light, compared to the usual derivation of the Dirac equation as *square root* of the Klein-Gordon equation.

Let us go back to the nonlinear realization approach of section 2. Apart from the connection (5), we have claimed that we can, for every field φ transforming under *some* representation of G , define the field $\psi = \sigma^{-1}\varphi$, that will transform under *some* representation of H . It is usually understood that, if φ transforms under a vector (spinor) representation of G , then ψ too will transform under a vector (spinor) representation of H . This means that σ itself will sometimes be interpreted as a mixed $G-H$ tensor, and sometimes as a mixed $G-H$ spinor. This is the usual conception in the nonlinear realization approach.

This may make sense, as long as one considers σ simply as transformation that parameterizes the degrees of freedom of the coset space G/H . However, as we have seen, in a dynamically broken theory, σ , represented by the reducing matrix r_a^α , will appear as the Nambu-Goldstone field of the theory. These fields, even though of pure gauge nature, describe the Nambu-Goldstone particles, and as such, have to transform under a well specified representation of G and of H . We have thus to decide whether r_a^α transforms as $H-G$ tensor (as was the case throughout our article) or, say, as $H-G$ spinor.

In this section, we are interested in the breakdown of the group $GL(R^4)$ to the Lorentz group. Since spinor representations of $GL(R^4)$ are necessarily infinite dimensional, we choose here a third way. We introduce the reducing matrix with the following transformation behavior (see (9)):

$$r_\alpha^M \rightarrow L_N^M r_\beta^N (G^{-1})^\beta_\alpha, \quad (103)$$

with $G^\alpha_\beta \in GL(R^4)$ and

$$L^M_N = e^{i\varepsilon^{ab}(\sigma_{ab})^M_N}, \quad (104)$$

where $\sigma_{ab} = (i/2)[\gamma_a, \gamma_b]$ are (twice) the generators of the spinor representation of the Lorentz group and $\varepsilon_{ab} \in R$. In other words, r^M_α transforms as a spinor with respect to the index M . More generally, from a $GL(R^4)$ vector k^α , we can now form the quantity

$$\psi^M = r^M_\alpha k^\alpha, \quad (105)$$

which transforms exactly like a Dirac spinor, $\psi \rightarrow e^{i\varepsilon^{ab}\sigma_{ab}}\psi$. Usually, this is called an $SL(2, C)$ transformation, and we will follow that tradition, although whenever we talk about spinors, we actually refer to 4-component Dirac spinors, or bispinors. Note also that we use capital latin letters from the middle of the alphabet to index the $SL(2, C)$ quantities, as opposed to $a, b, c, \dots$ which will be used, as before, for $SO(3, 1)$ indices.

We see that r^M_α is very similar to a spin 3/2 particle ψ_i , or ψ^M_i if we write the Dirac space index explicitly, with the only difference that the spacetime index i is replaced by a $GL(R^4)$ index. As we know, we can always choose a gauge in which we can identify those indices. Therefore, the corresponding Nambu-Goldstone particles will be of fermion nature.

An $SL(2, C)$ connection can also be defined using (8) and r^M_α . More physical than *reducing* the symmetry by hand is, of course, the dynamical Higgs approach, to which we will turn now.

As gravitational fields, we introduce a $GL(R^4)$ connection Γ^α_β and a tensor valued one-form e^α_β , whose relation to the tetrad field will be clarified in the following. In addition, instead of the metric $g_{\alpha\beta}$ of the previous sections, we consider a Higgs field γ^α_β with one co- and one contravariant index.

Before we construct the gravitational Lagrangian, let us point out some important differences between the metric g and the tensor γ . In contrast to the metric tensor, which transforms as $g \rightarrow G^{-1}g(G^{-1})^T$, the tensor γ will transform as $\gamma \rightarrow G\gamma G^{-1}$. Therefore, while the metric is entirely characterized by its signature, γ will be characterized by its four eigenvalues. These cannot be scaled to ± 1 . More generally, if we assign a groundstate to γ , there will always be a residual dilational invariance $G^\alpha_\beta = \lambda\delta^\alpha_\beta$ under which γ is invariant. On the other hand, in contrast to g , from γ we can form the $GL(R^4)$ invariants $\det \gamma$ and $Tr \gamma$.

Since γ contains a priori 16 independent components, while the broken gauge degrees of freedom are only 10, we are free to put constraints on γ . It turns out that it is

enough to require

$$\gamma^\alpha_\alpha = 0. \quad (106)$$

Let us define the gauge invariant spacetime metric as

$$g_{ik} = -e^\alpha_{\beta i} e^\beta_{\alpha k}. \quad (107)$$

Consider the following Higgs potential:

$$\sqrt{|g_{ik}|} V = \sqrt{|g_{ik}|} \left[\frac{1}{4} \text{Tr}(\gamma^4) + \frac{1}{2} \text{Tr}(\gamma^2) \right]. \quad (108)$$

The matrix notation is to be understood as $\text{Tr}(\gamma^2) = \gamma^\alpha_\beta \gamma^\beta_\alpha$ and similar for $\text{Tr}(\gamma^4)$. Variation with respect to γ leads to the groundstate characterization

$$\gamma^\alpha_\beta \gamma^\beta_\delta = -\delta^\alpha_\delta. \quad (109)$$

The other solution, $\gamma = 0$, corresponds to a maximum of V and is thus unstable. Although we did not explicitly mention it, it is understood that all the fields are real, since we are dealing with a $GL(R^4)$ theory.

The condition (109), together with the constraint (106), are enough to characterize γ completely. We see this as follows: Suppose we diagonalize γ . Then, (106) and (109) lead to the eigenvalues $(i, i, -i, -i)$. In other words, the eigenvalues are entirely fixed, and therefore γ itself is fixed up to similarity transformations. In our specific case, we cannot diagonalize the real matrix γ with a $GL(R^4)$ transformation, but this is also not necessary. We can choose any real representative of γ (with eigenvalues $(i, i, -i, -i)$), say

$$\gamma = \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix} = i \begin{pmatrix} \sigma^2 & 0 \\ 0 & -\sigma^2 \end{pmatrix} \quad (110)$$

and then, any other real solution to (106) and (109) will be related to this choice by a $GL(R^4)$ transformation.

The groundstate (110) certainly breaks the $GL(R^4)$ symmetry. But down to which subgroup? As mentioned earlier, the dilational symmetry cannot be broken. In order to find the rest of the residual symmetry group, it is convenient to introduce, at this

point, the following Dirac matrices

$$\begin{aligned}\gamma_0 &= \begin{pmatrix} 0 & \sigma^2 \\ \sigma^2 & 0 \end{pmatrix}, & \gamma_1 &= \begin{pmatrix} -i\sigma^3 & 0 \\ 0 & -i\sigma^3 \end{pmatrix}, \\ \gamma_2 &= \begin{pmatrix} 0 & \sigma^2 \\ -\sigma^2 & 0 \end{pmatrix}, & \gamma_3 &= \begin{pmatrix} i\sigma^1 & 0 \\ 0 & i\sigma^1 \end{pmatrix}.\end{aligned}\tag{111}$$

These are the usual Dirac matrices in the Majorana representation. We have the following relation:

$$\gamma_5 \equiv \gamma = \gamma_0 \gamma_1 \gamma_2 \gamma_3.\tag{112}$$

We attach the index 5 to our groundstate Higgs matrix, in order to indicate its relation to the Dirac matrices in the specific representation (and not to denote any kind of transformation behavior). Note that our γ_5 differs by a factor i from the usual definition.

Consider the exponential map, i.e., write the $GL(R^4)$ transformation (in the neighborhood of the identity I) in the form

$$G^\alpha_\beta = e^{\sigma^\alpha_\beta}\tag{113}$$

where σ some real 4×4 matrix. [Similarly, the elements in the neighborhood of $-I$ can be written as $-\exp(\sigma^\alpha_\beta)$. It is, however, evident that, if (110) is invariant under some transformation G , then also under $-G$.] As to the other connected component of $GL(R^4)$ (with $\det G < 0$), its elements can be written in the form

$$G^\alpha_\beta = \pm P e^{\sigma^\alpha_\beta},$$

with

$$P = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix},$$

the spatial inversion (or parity) transformation. It is easily checked that (110) is not invariant under P , and therefore, we can concentrate on the transformations of the form (113). The invariance of γ^α_β , i.e., the requirement $G\gamma_5 G^{-1} = \gamma_5$, leads infinitesimally to the following condition

$$\sigma\gamma_5 = \gamma_5\sigma.\tag{114}$$

It is now an easy task to determine the matrices σ that satisfy this condition. It is convenient to consider, temporarily, σ to be a complex matrix. We know from Dirac theory that the complete set of linearly independent (complex) matrices is given by the sixteen matrices $(I, \gamma_5, \gamma_a, \gamma_5\gamma_a, [\gamma_a, \gamma_b])$. Apart from I and γ_5 , corresponding to dilations and chiral transformations, to which we will turn later on, the only matrices satisfying (114) are the six matrices $[\gamma_a, \gamma_b]$. Therefore, the most general transformation allowed by (114) (up to dilations and chiral transformations), has the form

$$G = e^{\tilde{\varepsilon}^{ab}[\gamma_a, \gamma_b]}, \quad (115)$$

with, in general, complex parameters $\tilde{\varepsilon}^{ab}$. Let us introduce the Lorentz generators

$$\sigma_{ab} = \frac{i}{2}[\gamma_a, \gamma_b]. \quad (116)$$

More precisely, the matrices $\sigma_{ab}/2$ are the Lorentz generators, but it is customary to define σ_{ab} in the form (116). We see from (111) that all six σ_{ab} are purely imaginary. Therefore, since the exponent in (115) is real, we conclude that G has the form

$$G = e^{-\frac{i}{4}\varepsilon^{ab}\sigma_{ab}}, \quad (117)$$

with real parameters ε^{ab} . (The factor $-1/4$ is conventional.) This, however, is exactly the $SL(2, C)$ transformation (104). Note that the essential step was the reality requirement of the $GL(R^4)$ transformation. Without that, we can, in (115), replace any matrix γ_a by $i\gamma_a$, which is equivalent to a signature change of the metric $\eta_{ab} = \frac{1}{2}(\gamma_a\gamma_b + \gamma_b\gamma_a)$ and more generally to a change of the residual symmetry group. The reality requirement, on the other hand, allows only for a change $\gamma_a \rightarrow i\gamma_a$ of *all* matrices, corresponding to a global, irrelevant, sign change of the metric.

As before, we use the letters $M, N \dots$ to denote $SL(2, C)$ quantities. The requirement that the Dirac matrices $\gamma^a = (\gamma^a)_N^M$ are invariant under a gauge transformation leads to the usual relation between Lorentz and $SL(2, C)$ transformations, i.e.,

$$\gamma^a \rightarrow e^{-\frac{i}{4}\varepsilon^{cd}\sigma_{cd}}(\Lambda^a_b\gamma^b)e^{\frac{i}{4}\varepsilon^{cd}\sigma_{cd}} = \gamma^a, \quad (118)$$

which is satisfied (infinitesimally) for $\Lambda^a_b = \delta^a_b + \varepsilon^a_b$. [The same can be achieved replacing G by $-G$ in (117). This reflects the well known two-to-one correspondence between $SL(2, C)$ and $SO(3, 1)$.] After the symmetry breakdown, the $GL(R^4)$ connection Γ^α_β reduces to a $SL(2, C)$ connection Γ_N^M and the $GL(R^4)$ tensor e^α_β to the $SL(2, C)$ tensor e_N^M . Following Chamseddine [31], we can introduce $SO(3, 1)$ components in the

following way

$$\begin{aligned} e_N^M &= ie^a(\gamma_a)_N^M, \\ \Gamma_N^M &= -\frac{i}{4}\Gamma^{ab}(\sigma_{ab})_N^M. \end{aligned} \quad (119)$$

Since Γ_N^M and e_N^M are real, the same holds for the Lorentz connection Γ^{ab} and the tetrad field e^a . The inverse relations are

$$\begin{aligned} e^a &= -\frac{i}{4} \text{Tr} (e \gamma^a), \\ \Gamma^{[ab]} &= \frac{i}{2} \text{Tr} (\Gamma \sigma^{ab}), \end{aligned} \quad (120)$$

where the traces are to be taken in Dirac space, and $e = e_N^M$, $\Gamma = \Gamma_N^M$. We see that (119) parameterizes only the antisymmetric part of the connection Γ^{ab} . The non-metricity parts are lost. We will return to this point later on. Note also that the spacetime metric (107) takes the expected form $g_{ik} = e_i^a e_k^b \eta_{ab}$.

Let us now construct the Lagrangian of the theory. Consider the following $GL(R^4)$ invariant Lagrangian:

$$\mathcal{L}_0 = \frac{1}{4} \gamma^\alpha_\beta R^\beta_\gamma \wedge e^\gamma_\delta \wedge e^\delta_\alpha. \quad (121)$$

In the groundstate $\gamma = \gamma_5$, this simply reduces to

$$\mathcal{L}_0 = \frac{1}{4} \text{Tr} [\gamma_5 R \wedge e \wedge e], \quad (122)$$

where we use the matrix notation, as is customary in Dirac space. This, however, is exactly Chamseddine's form of the $SL(2, C)$ invariant Einstein-Cartan Lagrangian [31]. Indeed, using (119) and (120), we find $R_N^M = -(i/2)R^{ab}(\sigma_{ab})_N^M$ and

$$\mathcal{L}_0 = \frac{1}{2} \varepsilon_{abcd} R^{ab} \wedge e^c \wedge e^d, \quad (123)$$

or, in the conventional form, $\mathcal{L}_0 = -2eR$.

Until this point, nothing seems to have been gained compared to the usual metric approach of the affine theory. On the contrary, several issues are still open, like the discussion of the dilations and chiral transformations, as well as the parameterization of the non-metricity. We postpone the full analysis of those problems to future work, although some insight will also be gained in the following.

However, when it comes to spinor fields, the approach of this section clearly presents undeniable advantages. This is the point we concentrate on in this article.

The presence of the tensor valued one-form e^α_β allows us to write down a Lagrangian for a vector field k^α of first order in the derivatives. More precisely, consider the two vector fields k^α and k_α . (Vectors under $GL(R^4)$, scalars under spacetime transformations.) Since we are not in the possession of a $GL(R^4)$ metric that could relate a covariant to a contravariant vector, these fields have to be considered as independent from each other. Moreover, we allow both fields to take complex values, with the only restriction $k_\alpha k^\alpha \in R$.

Consider the Lagrangian

$$\begin{aligned} \mathcal{L}_M = & \frac{1}{48} [k_\alpha e^\alpha_\beta \wedge e^\beta_\gamma \wedge e^\gamma_\delta \wedge \gamma^\delta_\epsilon Dk^\epsilon - Dk_\alpha \wedge e^\alpha_\beta \wedge e^\beta_\gamma \wedge e^\gamma_\delta \gamma^\delta_\epsilon k^\epsilon] \\ & + \sqrt{|\det g|} m k_\gamma k^\gamma, \end{aligned} \quad (124)$$

with $Dk_\alpha = dk_\alpha - \Gamma^\beta_\alpha k_\beta$ and $Dk^\alpha = dk^\alpha + \Gamma^\alpha_\beta k^\beta$. We claim that (124) is the Dirac Lagrangian in a general linear invariant form. Indeed, in the groundstate $\gamma = \gamma_5$, using the parameterizations (119) and changing, as in (105), the notation to $k^M = \psi^M$ and $k_M = \bar{\psi}_M$, (124) reduces to

$$\begin{aligned} \mathcal{L}_M = & -\frac{i}{12} \left[\bar{\psi} (e^a \wedge e^b \wedge e^c \varepsilon_{abcd} \gamma^d) \wedge D\psi - \frac{i}{12} D\bar{\psi} \wedge (e^a \wedge e^b \wedge e^c \varepsilon_{abcd} \gamma^d) \psi \right] \\ & + e m \bar{\psi} \psi \\ = & e \left(\frac{i}{2} [\bar{\psi} \gamma^m D_m \psi - (D_m \bar{\psi}) \gamma^m \psi] + m \bar{\psi} \psi \right) \end{aligned} \quad (125)$$

with $D\psi = d\psi - \frac{i}{4} \Gamma^{ab} \sigma_{ab} \psi$, $D\bar{\psi} = d\bar{\psi} + \frac{i}{4} \bar{\psi} \Gamma^{ab} \sigma_{ab}$, $\gamma^m = e^m_a \gamma^a$ and $e = \det e^a_m$. This is the well known form of the Lorentz invariant Dirac Lagrangian in the framework of Poincaré gauge theory [20].

The complete Lagrangian will be composed from (108), (121) and (124), and possibly a kinetic term for the Higgs field γ .

It is easy to check that $\psi, \bar{\psi}, D\psi$ and $D\bar{\psi}$ have the required $SL(2, C)$ transformation behavior known from Dirac theory. This is evident anyway, by construction. There remains one point to discuss, namely the relation between k^α and k_α , or ψ and $\bar{\psi}$ respectively.

We have argued that k_α and k^α have to be considered as independent, due to the lack of a $GL(R^4)$ metric. However, after the symmetry breakdown, the residual symmetry is $SL(2, C)$ (neglecting again the dilations and chiral transformations). Similar to the Minkowski metric in the Lorentz case, one might look for a non-dynamical, constant tensor that is invariant under $SL(2, C)$. In other words, is there an invariant tensor g_{MN} that could relate covariant and contravariant $SL(2, C)$ vectors in the form (take care of the index positions, g_{NM} may be asymmetric)

$$k_M = g_{NM} k^N? \quad (126)$$

Note that, as opposed to the γ^a matrices, g_{MN} has two lower indices and therefore transforms as

$$g \rightarrow e^{-i\varepsilon^{ab}\sigma_{ab}} g (e^{-i\varepsilon^{ab}\sigma_{ab}})^T. \quad (127)$$

Infinitesimally, the requirement $g \rightarrow g$ takes the form

$$\sigma g + g \sigma^T = 0, \quad (128)$$

with σ one of the matrices σ_{ab} . Considering again the 16 linearly independent 4x4 matrices, we find that there is only one matrix that anti-commutes in this way with all the matrices σ_{ab} , namely

$$g_{MN} = \alpha (\gamma_0)_{MN}. \quad (129)$$

Note, by the way, that γ_5 does not anti-commute in this way with $g_{MN} = (\gamma_0)_{MN}$, and therefore, the requirement of the existence of the $SL(2, C)$ invariant metric also breaks the chiral invariance. Finally, the requirement that g_{MN} is a non-dynamical, constant matrix (i.e., $\alpha = \text{const}$), also breaks the dilational gauge freedom. We may choose $\alpha = 1$. Let us emphasize that the matrix $(\gamma_0)_{MN}$ in (129) has nothing to do with the Dirac matrix $(\gamma_0)_N^M$. Apart from the different index positions, there is also the important difference that in (129), the index 0 does not mean that the matrix is a component of a Lorentz vector (as is the case with $(\gamma_0)_N^M$), but it is attached only to indicate the (rather incidental) numerical coincidence of both matrices.

However, the relation $k_M = (\gamma_0)_{NM} k^N$ is easily shown to lead to $k_M k^M = 0$ for any $SL(2, C)$ vector k^M . Therefore, we replace (126) by

$$k_M = g_{NM} (k^N)^*, \quad \text{with } g_{NM} = (\gamma_0)_{NM}, \quad (130)$$

where the star denotes complex conjugation. Note that, for real k^M , we are lead to $k^M k_M = 0$. This is the reason why it is necessary to introduce complex fields. [Recall in this context that, in the Majorana representation, the charge conjugation operator

C is given by $-\gamma^0$ and therefore $\psi_c = \psi^*$.] In matrix notation, and writing again $k^M = \psi^M, k_M = \bar{\psi}_M$, (130) reads

$$\bar{\psi} = \psi^t \gamma_0, \quad (131)$$

with $\psi^t = (\psi^*)^T$ the hermitian conjugate. This is the well known relation for the adjoint spinor in Dirac theory. Note that (130) assures that $k_M k^M$, i.e., $\bar{\psi}\psi$, is automatically real, which justifies the choice of $\alpha = 1$ in (129).

We conclude that the requirement of the existence of an $SL(2, C)$ invariant matrix, that plays the role of a metric tensor in Dirac space, in the sense of equation (130), leads to the usual relation between the spinor and the adjoint spinor field. The introduction of this tensor, which is numerically equal to γ_0 , also breaks the dilational and chiral invariance that were still symmetries of the groundstate (110).

The metric (129) is also useful under another aspect. It can be used to raise the second index of the $SL(2, C)$ connection Γ_N^M , which allows us to consider the symmetric and antisymmetric parts of Γ^{MN} . It is then easy to show that the antisymmetric part transforms as tensor under $SL(2, C)$, whereas the symmetric part has a non-homogeneous transformation behavior. The situation is thus reversed compared to the Lorentz connection Γ^{ab} . It also becomes clear that the parameterization (119) presupposes that Γ^{MN} and e^{MN} are symmetric, because the matrices σ_{MN} and $(\gamma^a)_{MN}$ (where one index has been lowered with $g_{MN} = (\gamma_0)_{MN}$) are symmetric.

Therefore, in order to parameterize completely the connection Γ_N^M , one will have to use the sixteen linearly independent matrices in Dirac space and write, instead of (119),

$$\Gamma_N^M = -\frac{i}{4} \Gamma^{ab} (\sigma_{ab})_N^M + iA^a (\gamma_a)_N^M + iB^a (\gamma_a \gamma_5)_N^M + C \delta_N^M + D (\gamma_5)_N^M, \quad (132)$$

and in the following, analyze the transformation behavior of the one-forms A^a, B^a, C, D , as well as their coupling to the Dirac field. This parameterization of the $GL(R^4)$ connection in terms of the irreducible components under the Lorentz group is rather similar to the case of the $SO(4, 2)$ connection (conformal group) (see [16] for instance), the difference lying in the dilational part $C \delta_N^M$ which is not present in the conformal case. One will also have to put certain constraints on e^α_β , because else, e_N^M will allow for a similar parameterization, and the direct correspondence to e^a will be lost.

In our quest for a minimalistic model, there is also the attractive idea to consider the tetrad as being related to the (pseudo)vector part A^a or B^a of the connection (132). Indeed, with (132), the curvature tensor (after the symmetry breaking) takes

the following form

$$\begin{aligned}
R &= d\Gamma + \Gamma \wedge \Gamma \\
&= -\frac{i}{4}R^{ab}\sigma_{ab} + i(A^a \wedge A^b + B^a \wedge B^b)\sigma_{ab} + i(dA^a + \Gamma^a_c \wedge A^c)\gamma_a \\
&\quad - 2i(B^a \wedge D)\gamma_a + i(dB^a + \Gamma^a_c \wedge B^c)\gamma_a\gamma_5 + 2i(A^a \wedge D)\gamma_a\gamma_5 \\
&\quad + dD\gamma_5 - 2(A^a \wedge B_a)\gamma_5 + dC.
\end{aligned} \tag{133}$$

Therefore, one could consider, instead of (121) the following Lagrangian

$$\begin{aligned}
\mathcal{L}_0 &= Tr(\gamma R \wedge R) \\
&= -4(dD - 2A^a \wedge B_a) \wedge dC \\
&\quad - 4\varepsilon_{abcd}[\frac{1}{4}R^{ab} - A^a \wedge A^b - B^a \wedge B^b] \\
&\quad \wedge [\frac{1}{4}R^{cd} - A^c \wedge A^d - B^c \wedge B^d].
\end{aligned} \tag{134}$$

The first line is the $GL(R^4)$ invariant form, depending only on Γ^α_β and the Higgs field γ^α_β , and the following lines are its decomposition in the groundstate $\gamma = \gamma_5$. Under the assumption that the pseudo-vector part B^a vanishes, we can identify $A^a \sim e^a$, and (134) reduces to the Einstein-Cartan Lagrangian with cosmological constant. (The terms $dC \wedge dD$ and $\varepsilon_{abcd}R^{ab} \wedge R^{cd}$ are total divergences.)

In order to write down the Dirac Lagrangian (124), we define the tensor valued one-form

$$e \equiv \frac{1}{2}\gamma D\gamma = \frac{1}{2}\gamma d\gamma + \frac{1}{2}\gamma[\Gamma, \gamma], \tag{135}$$

which decomposes in the groundstate to

$$e = iA^a\gamma_a + iB^a\gamma_a\gamma_5. \tag{136}$$

Again under the assumption $B^a = 0$, this reduces to the the form of e given in (119), and can be used in (124), as well as in (107),(108). Of course, one will have to justify the assumption $B^a = 0$. Another possibility would be to consider left and right handed tetrad fields ($A^a = -B^a$ and $A^a = B^a$ in (136)) and somehow to exclude one of both polarizations.

We see that, in this way, one can write down a gravitational theory based on $GL(R^4)$ only, without gauging the translational group.

8 Conclusions

We have constructed the Higgs sector of gauge theories of gravity using a minimum of additional structures, and presented the symmetry breaking from the original symmetry group down to the Lorentz stability subgroup, following an approach suggested by the nonlinear realization treatment of those groups. The results of Stelle and West [11] have been reproduced and the analysis has been extended to the Poincaré group, the affine group and the group $SL(R^5)$.

In the case of the Poincaré group, the following conclusions have been drawn:

- 1) No Poincaré Lagrangian that leads to a nontrivial Einstein equation can be written down without the help of an additional Poincaré vector y^a .
- 2) In order to assign a physical meaning to y^a , it has to be interpreted as Higgs field that triggers the symmetry breaking of the translational gauge freedom.
- 3) No Poincaré invariant Higgs potential can be written down. This is also not necessary, since the mere presence of a Poincaré vector automatically breaks the symmetry if some groundstate value $y^a(0)$ is assigned to it.
- 4) The most general kinetic term for y^a takes the form of a cosmological constant. The *natural* sign of the kinetic term favors the choice of a positive cosmological constant. A negative or zero value, however, is not excluded on theoretical grounds.

The following conclusions result from the analysis of the affine symmetry group:

- 1) As in point 1 and 2 above, the general linear metric $g_{\alpha\beta}$, as well as the affine vector y^α are necessary to write down a meaningful Lagrangian for the gravitational fields and they should be treated as Higgs fields.
- 2) The translational part of the affine group is treated in exactly the same way as in the case of the Poincaré group. Consequently, conclusions 3 and 4 above are valid also in this case.
- 3) The symmetry breakdown of the special linear group SL down to the Lorentz group is easily triggered by a Higgs sector that assigns a negative value to the metric determinant in the groundstate. This fixes the metric signature to the Lorentzian one and therefore fixes the stability subgroup to $O(3,1)$. The same signature is passed on to the physical spacetime metric. Matter fields that couple directly to the symmetric

part of the GL connection (i.e., possessing hypermomentum) may provoke a signature change.

4) In order to generalize the above procedure to the general linear group GL , the introduction of an additional *dilaton* field, a scalar density of weight $+1$, cannot be avoided. The reason for this is traced back to the fact that, in metric affine theory, one of the equations obtained from the variation with respect to the tetrad and the metric respectively, is always redundant. This has its origin in the pure gauge nature of the tetrad field, which can be gauged into the non dynamical form δ_i^α by a GL transformation. As a result, we cannot get an independent equation for $g_{\alpha\beta}$ that would allow us to fix the groundstate of the theory. This problem can only be resolved by the introduction of an additional matter field, the dilaton φ , which is conveniently chosen to transform as scalar density, and provides us with an additional, independent equation.

5) The introduction of a Higgs sector for the dilaton field, initially intended to break down the GL group to SL , in an approach similar to that of Hehl et al. [20], reveals itself as not enough to assign a groundstate value to φ . Only under the assumption that φ is nonzero, which implicitly means that we break the symmetry down to SL , the equations determine the sign of the metric determinant, and thus, lead to the final, Lorentz invariant groundstate metric. In other words, the dilaton field φ allows us to use the same procedure, as described under point 3, in the framework of the general linear group, without however assigning a nonzero value to φ , which has to be assumed ad hoc.

6) We were able to present a solution to this problem, by introducing, in addition to the dilaton, a scalar field ψ . In this way, we get a complete symmetry breaking from GL down to $O(3,1)$, without any further assumptions. In the absence of matter fields with hypermomentum, the signature will remain Lorentzian, and moreover, the Higgs sector will not contribute to the remaining gravitational equations, which means that the theory essentially reduces to Poincaré gauge theory.

7) A similar approach cannot be used to construct a theory with a residual $O(4)$ or $O(2,2)$ symmetry, since the knowledge of the metric determinant in those cases will not uniquely specify the stability subgroup. The Higgs sector of such theories would have to be constructed directly in terms of the signature (as a GL invariant), which leads to highly nontrivial expressions. Thus, the Lorentz group, as opposed to the other rotation groups, seems to be favored by nature.

As to the group $SL(R^5)$, we concluded the following:

- 1) The $SL(R^5)$ tensor g_{AB} and vector y^A are enough to trigger the breakdown of the symmetry group down to the Lorentz group. No additional structure is needed.
- 2) Comparing with the affine case, the dilaton field appears now as the g_{55} component of the metric tensor and the scalar field can be found in the quantity $g_{AB}y^Ay^B$. A natural origin of those fields, introduced ad hoc in the affine theory, is therefore provided. Roughly, to the affine Higgs fields $g_{\alpha\beta}, y^\alpha, \varphi, \psi$ correspond the $SL(R^5)$ Higgs fields $g_{\alpha\beta}, y^\alpha, g_{55}, y^5$, whereas the remaining fields, namely $g_{\alpha 5}$, are responsible for the breakdown of the additional, pseudo-inversional symmetry.

As a final conclusion, we see that the affine group as symmetry group of the gravitational interaction presents some interesting features that are not present in other gravitational gauge theories. These are its direct relation to general relativity and to metric theories in general, and the possibility of explaining dynamically the signature of physical spacetime. However, the symmetry breaking mechanism, in the form presented in this article, takes a more natural form in the framework of the theory based on the special linear group $SL(R^5)$. Therefore, it seems promising to take a closer look at that theory in future work.

An alternative to the metric based models has been presented in the last section. It has been shown that a $GL(R^4)$ covariant tensor, which turns out to be related to the Dirac matrix γ_5 , can be used to break the general linear group down to the group $SL(2, C)$, which is isomorphic to the Lorentz group. This model, although still incomplete in many aspects, seems especially promising as far as the incorporation of spinor fields is concerned. A general linear covariant generalization of the Dirac equation can easily be written down using only finite dimensional vector representations.

References

- [1] J. Goldstone, Nuovo Cimento **19**, 154 (1961)
- [2] P.W. Higgs, Phys. Lett. **12**, 132 (1964)
- [3] Y. Nambu, Phys. Rev. Lett. **4**, 380 (1960)
- [4] S. Weinberg, Phys. Rev. Lett. **19**, 1254 (1967)

- [5] S. Coleman, J. Wess, and B. Zumino, Phys. Rev. **177**, 2239 (1969); C.G. Callan, S. Coleman, J. Wess, and B. Zumino, Phys. Rev. **177**, 2247 (1969)
- [6] A. Salam and J. Strathdee, Phys. Rev. **184**, 1750 (1969); **184**, 1760 (1969)
- [7] C.J. Isham, A. Salam, and J. Strathdee, Ann. Phys. (N.Y.) **62**, 98 (1971)
- [8] L.N. Chang and F. Mansouri, Phys. Rev. D **17**, 3168 (1978)
- [9] F. Gürsey and L. Marchildon, Phys. Rev. D **17**, 2038 (1978)
- [10] A.A. Tseytlin, Phys. Rev. D **26**, 3327 (1982)
- [11] K.S. Stelle and P.C. West, Phys. Rev. D **21**, 1466 (1980)
- [12] Y. Ne'eman and D. Sijacki, Phys. Lett. B **200**, 489 (1988)
- [13] E.A. Lord, J. Math. Phys. **27**, 2415 (1986); **27**, 3051 (1986)
- [14] E.A. Lord and P. Goswami, J. Math. Phys. **29**, 258 (1988)
- [15] E.A. Ivanov and J. Niederle, Phys. Rev. D **25**, 976 (1982); **25**, 988 (1982)
- [16] J. Julve, A. Lopez-Pinto, A. Tiemblo, and R. Tresguerres, Class. Quantum Grav. **12**, 1327 (1995)
- [17] G. Grignani and G. Nardelli, Phys. Rev. D **45**, 2719 (1992)
- [18] E.W. Mielke, J.D. McCrea, Y. Ne'eman, and F.W. Hehl, Phys. Rev. D **48**, 673 (1993)
- [19] R. Tresguerres and E.W. Mielke, Phys. Rev. D **62**, 044004 (2000)
- [20] F.W. Hehl, J.D. McCrea, E.W. Mielke, and Y. Ne'eman, Phys. Rep. **258**, 1 (1995)
- [21] I. Kirsch and D. Sijacki, Class. Quantum Grav. **19**, 3157 (2002)
- [22] D. Ivanenko and G. Sardanashvily, Phys. Rept. **94**, 1 (1983)
- [23] G.A. Sardanashvily, Theor. Math. Phys. **132**, 1163 (2002)
- [24] G. Sardanashvily, J. Math. Phys. **39**, 4874 (1998)
- [25] A.B. Borisov and V.I. Ogievetsky, Theor. Math. Phys. **21**, 1179 (1975)

- [26] K. Hayashi, K. Nomura, and T. Shirafuji, Prog. Theor. Phys. **56**, 1239 (1991)
- [27] R. Tresguerres, Phys. Rev. D **66**, 064025 (2002)
- [28] J. Greensite, Phys. Lett. B **300**, 34 (1993)
- [29] S.D. Odintsov, A. Romeo, and R.W. Tucker, Class. Quant. Grav. **11**, 2951 (1994)
- [30] E.A. Ivanov and V.I. Ogievetsky, Teor. Mat. Fiz. **25**, 164 (1975)
- [31] A. H. Chamseddine, Phys. Rev. D **69**, 024015 (2004)